

# On a Potential Global Role for Vitamin K-dependent $\gamma$ -Carboxylation in Animal Systems

EVIDENCE FOR A  $\gamma$ -GLUTAMYL CARBOXYLASE IN *DROSOPHILA*\*

Received for publication, October 19, 2000  
Published, JBC Papers in Press, December 7, 2000, DOI 10.1074/jbc.M009576200

Craig S. Walker<sup>‡</sup>, Reshma P. Shetty<sup>‡§</sup>, Kathleen Clark<sup>‡</sup>, Sandra G. Kazuko<sup>¶</sup>, Anthea Letsou<sup>¶</sup>,  
Baldomero M. Olivera<sup>‡</sup>, and Pradip K. Bandyopadhyay<sup>‡¶</sup>

From the Departments of <sup>‡</sup>Biology and <sup>¶</sup>Human Genetics, University of Utah, Salt Lake City, Utah 84112-0840

The vitamin K-dependent  $\gamma$ -carboxylation of glutamate to  $\gamma$ -carboxyglutamate was originally well characterized in the mammalian blood clotting cascade.  $\gamma$ -Carboxyglutamate has also been found in a number of other mammalian proteins and in neuropeptides from the venoms of marine snails belonging to the genus *Conus*, suggesting wider prevalence of  $\gamma$ -carboxylation. We demonstrate that an open reading frame from a *Drosophila melanogaster* cDNA clone encodes a protein with vitamin K-dependent  $\gamma$ -carboxylase activity. The open reading frame, 670 amino acids in length, is truncated at the C-terminal end compared with mammalian  $\gamma$ -carboxylase, which is 758 amino acids. The mammalian gene has 14 introns; in *Drosophila* there are two much shorter introns but in positions precisely homologous to two of the mammalian introns. In addition, a deletion of 6 nucleotides is observed when cDNA and genomic sequences are compared. *In situ* hybridization to fixed embryos indicated ubiquitous presence of carboxylase mRNA throughout embryogenesis. Northern blot analysis revealed increased mRNA levels in 12–24-h embryos. The continued presence of carboxylase mRNA suggests that it plays an important role during embryogenesis. Although the model substrate FLEEL is carboxylated by the enzyme, a substrate containing the propeptide of a *Conus* carboxylase substrate, conantokin G, is poorly carboxylated. Its occurrence in vertebrates, molluscan systems (*i.e.* *Conus*), and *Drosophila* and the apparently strong homology between the three systems suggest that this is a highly conserved and widely distributed post-translational modification in biological systems.

The functions of proteins are coordinated physiologically by post-translational modification. For example, phosphorylation-dephosphorylation cascades integrate the biochemistry of individual proteins into cellular physiology. In addition to post-translational modifications that occur primarily within cells,

post-translational modifications also occur on extracellular proteins. The most familiar of these are *N*-glycosylation of asparagine residues and *O*-glycosylation of serine and threonine residues.

One of the most distinctive of the extracellular post-translational modifications is the vitamin K-dependent  $\gamma$ -carboxylation of glutamate residues to give  $\gamma$ -carboxyglutamate (1). When it was first characterized,  $\gamma$ -carboxylation was thought to be a biochemical specialization of the mammalian blood-clotting cascade. However, several bone proteins (2, 3) as well as an extracellular ligand, gas6 (4), were subsequently identified as having the post-translational modification, although in the latter cases the precise mechanistic role of  $\gamma$ -carboxylation for proper protein function has not been established definitively. In addition, two novel proline-rich  $\gamma$ -carboxyglutamic acid-containing proteins, PRGP1 and PRGP2, of unknown function have been identified (5).

Long after its characterization in blood-clotting factors, vitamin K-dependent  $\gamma$ -carboxylation of glutamate residues was discovered in a phylogenetically distant system: the neuropeptides made in the venom duct of the predatory cone snails *Conus* (6, 7). The venoms of these snails have ~100 different peptides; ~5% of these are believed to be  $\gamma$ -carboxylated (8). This post-translational modification has been found in a number of diverse *Conus* peptides but has been studied most intensively in an unusual *Conus* neuropeptide family, the conantokins, which are NMDA receptor antagonists.

In the conantokins, the significance of the post-translational modification can readily be demonstrated: these peptides are inactive in analogs without  $\gamma$ -carboxylation of glutamate residues. Incomplete  $\gamma$ -carboxylation of blood-clotting factors results in poor coagulation. It has been postulated that  $\gamma$ -carboxylation of both the conantokins and of factors of the blood-clotting cascade induces a helical conformation in the post-translationally modified regions. This postulated role of  $\gamma$ -carboxylation in determining conantokin structure has been generally supported by a number of subsequent structural studies on various conantokins (9–12).  $\gamma$ -Carboxyglutamic acid confers the property of  $\text{Ca}^{2+}$  binding to the modified protein. In the case of the blood-clotting factors, the binding to  $\text{Ca}^{2+}$  results in a conformational change exposing hydrophobic residues for interaction with membranes (13–17).

The enzymatic reaction in the invertebrate system has recently been shown to have many striking similarities (*e.g.* a requirement for reduced vitamin K and the presence of a  $\gamma$ -carboxylation recognition site on the substrate) to that of the  $\gamma$ -carboxylation of factors involved in the mammalian blood-clotting cascade (18, 19). Despite the clear functional importance of  $\gamma$ -carboxylation in these two disparate phylogenetic systems,  $\gamma$ -carboxylation of glutamate residues has been re-

\* This work was supported in part by a grant from the University of Utah Research Foundation (to P. K. B.), the Funding Initiative Seed Grant Program (1999), National Institutes of Health Grant GM48677, and Cognetix, Inc. (Salt Lake City, UT). The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

The nucleotide sequence(s) reported in this paper has been submitted to the GenBank™/EBI Data Bank with accession number(s) AF170280.

§ Supported by a Pfizer summer undergraduate research fellowship and a Beckman undergraduate research fellowship.

¶ To whom correspondence should be addressed: Dept. of Biology, University of Utah, 257 S. 1400 E., Rm. 201, Salt Lake City, UT 84112-0840. Tel.: 801-581-5907; Fax: 801-585-5010; E-mail: bandyop@biology.utah.edu.

garded as a highly specialized post-translational modification. In this report, we provide evidence that is strongly consistent with vitamin K-dependent  $\gamma$ -carboxylation in fact being a much more widely distributed biological phenomenon. We demonstrate by molecular techniques the presence of a vitamin K-dependent  $\gamma$ -carboxylase-related protein that is expressed in the fruit fly *Drosophila melanogaster*, which has a high degree of sequence identity with the mammalian enzyme. Similar observations have recently been reported by Li *et al.* (20). Although the role of  $\gamma$ -carboxylation in *Drosophila* remains unknown, this post-translational modification is present in arthropods, suggesting that it is generally distributed in animal systems. The strong conservation in sequence of the  $\gamma$ -glutamyl carboxylase in *Drosophila* and in mammals suggests an important functional role for the enzyme, resulting in strong selection for sequence conservation.

#### EXPERIMENTAL PROCEDURES

**Materials**—*Conus textile* venom ducts were obtained from Dr. L. J. Cruz (University of the Philippines). Vitamin K (phytonadione) was from Abbott Laboratories, and  $\text{NaH}^{14}\text{CO}_3$  (55 mCi/mmol) was from PerkinElmer Life Sciences. Enzymes were purchased from Life Technologies, Inc. PCR<sup>1</sup> reactions were performed in an Air Thermo-Cycler (Idaho Technology). Oligonucleotides were synthesized at the peptide sequencing facility at the University of Utah.

**Preparation of mRNA**—Adult *Drosophila* (Oregon) were frozen in liquid nitrogen and ground to a fine powder, and total RNA was isolated (21). Poly(A)<sup>+</sup> RNA was isolated using a Qiagen Oligotex mRNA kit according to the vendor's instructions. Molecular biology experiments were done according to methods described by Sambrook *et al.* (22).

**Sequence Analysis**—The cDNA sequence of *Drosophila*  $\gamma$ -carboxylase was assembled from sequences of PCR products obtained by amplification of oligo(dT)- or Q<sub>17</sub>-primed cDNA using primers shown in Table I. The primer combinations used in the PCRs are shown in Table I. Primers 1 and 2 correspond to amino acid sequences conserved in human, bovine, and rat. Primers 5 and 6 were selected from the *Drosophila* genomic sequence (Berkeley *Drosophila* Genome Project, accession number AC005557). 3'-Sequences of the carboxylase mRNA were determined by the technique of rapid amplification of cDNA ends described by Frohman (23). The PCR product was cloned into the TA cloning vector (Invitrogen), and the nucleic acid sequence was determined. DNA sequencing was performed using ABI Prism BigDye terminators and cycle sequencing with *Taq* FS DNA polymerase (Life Technologies). The DNA sequence was collected and analyzed on an ABI Prism 377 automated DNA sequencer (Applied Biosystems, Foster City, CA). To obtain the 5' end of the coding sequences, we selected primer 6 from the genomic sequence. This allowed us to characterize a transcript with an open reading frame from nucleotides 52 (ATG) to 2198 (TGA). The sequences have been submitted to GenBank; the accession number is AF170280.

Analyses of amino acid homology between human (24), bovine (25), rat (26), and *Drosophila*  $\gamma$ -glutamyl carboxylases were carried out using Gap and PileUp programs version 4.0, 1998 (Genetics Computer Group).

**Northern Blot Analysis**—Total RNA was isolated from *Drosophila* at various stages of development. Embryos at 0–2, 2–4, 4–8, 12–16, and 16–24 h of development, larval stages 1–3, pupae, and adult flies were used in the experiment. Northern blot analysis was performed using reagents provided in the Northern Max kit (Ambion). Nine  $\mu\text{g}$  of total RNA from each sample were electrophoresed in a 1.5% denaturing agarose gel, transferred to a Gene Screen Plus membrane (PerkinElmer Life Sciences), and hybridized to [ $\alpha$ -<sup>32</sup>P]UTP-labeled antisense *Drosophila*  $\gamma$ -carboxylase RNA. As a control for loading, a membrane containing identical samples was hybridized to [ $\alpha$ -<sup>32</sup>P]dCTP-labeled *Drosophila* rp49. RNA molecular weight standards (RNA Millennium) were purchased from Ambion. Membranes were exposed to Molecular Dynamics (Sunnyvale, CA) Phosphor Screen and scanned. Images were analyzed using the NIH Image processing program.

**In Situ Hybridization**—The spatial distribution of  $\gamma$ -glutamyl carboxylase RNA was probed by hybridization to whole-mount embryos *in*




FIG. 1. Vectors used for the expression of GFP,  $\gamma$ -carboxylase-GFP, and  $\gamma$ -carboxylase in S2 cells.

*situ* (27). The cDNA was cloned into a dual-promoter (T7 and Sp6) vector. Both sense and antisense RNAs were synthesized using the appropriate RNA polymerase in the presence of digoxigenin-labeled uridine triphosphate. The digoxigenin-labeled RNA was used as probe in hybridizations to fixed embryos. The hybridized digoxigenin-labeled RNA was detected by incubating the embryos with alkaline phosphatase-conjugated anti-digoxigenin antibody and developed using nitroblue tetrazolium and 5-bromo-4-chloro-3-indolyl phosphate.

**Transient Expression of  $\gamma$ -Glutamyl Carboxylase Activity**—DNA containing the coding sequence of *Drosophila*  $\gamma$ -glutamyl carboxylase was obtained by PCR amplification of the genomic sequences. The 3' PCR primer was designed such that the  $\gamma$ -carboxylase (GC) coding sequences would be in frame with coding sequences of green fluorescent protein (GFP) in the expression plasmid pRmHa-3.GFP (Fig. 1). The  $\gamma$ -glutamyl carboxylase encoding sequences include the two introns in the genomic sequence. pRmHa-3.GFP was constructed by introducing the coding sequence of GFP from the pEGFP vector (CLONTECH) into pRmHa-3 (28). Expression in this plasmid is under the control of the inducible metallothionein promoter and carries the alcohol dehydrogenase poly(A) addition signal. *Drosophila* Schneider 2 (S2) cells were transfected with pRmHa-3.GC.GFP DNA using CellFECTIN (Life Technologies). 24 h after transfection cells were induced with 0.7 mM  $\text{CuSO}_4$ . 48 h after induction 50% of the cells expressed GFP as judged by fluorescent microscopy. The results also indicated that the introns were properly processed, and a continuous reading frame was present in the cloned GC. pRmHa-3.GC.GFP was modified to introduce a stop codon at the end of the GC coding sequences and to delete the GFP coding sequences. The modified plasmid pRmHa-3.GC\* was transfected into S2 cells. The cells were induced with 0.7 mM  $\text{CuSO}_4$  and harvested 48 h after induction. Cells were washed twice with phosphate-buffered saline and resuspended in buffer containing 25 mM 4-morpholinepropanesulfonic acid, pH 7.0, 0.5 M NaCl, 0.2% 3-[(3-cholamidopropyl)dime-thylammonio]-1-propanesulfonic acid/phosphatidyl choline, 2 mM EDTA, 2 mM dithiothreitol, 0.2  $\mu\text{g}/\text{ml}$  leupeptin, 0.8  $\mu\text{g}/\text{ml}$  pepstatin, and 0.04 mg/ml phenylmethylsulfonyl fluoride. The cell suspension was briefly sonicated using a Branson 450 sonifier and incubated in ice for 20 min. The lysate was assayed for carboxylase activity.

**Enzyme Assays**— $\gamma$ -Glutamyl carboxylase assays were performed as described by Stanley *et al.* (18). Reactions were done in a total volume of 125  $\mu\text{l}$  containing cell lysate and a final concentration of reagents as follows: 25 mM 4-morpholinepropanesulfonic acid, pH 7.4, 0.5 M NaCl, 0.2% 3-[(3-cholamidopropyl)dime-thylammonio]-1-propanesulfonic acid, 0.2% phosphatidyl choline, 0.8 M ammonium sulfate, 5  $\mu\text{Ci}$  of  $\text{NaH}^{14}\text{CO}_3$ , 6 mM dithiothreitol, 222  $\mu\text{M}$  reduced vitamin K, and 1.2 mM of a model substrate, the pentapeptide FLEEL (29). Reaction mixtures were incubated at 25 °C for 120 min and were quenched by addition of 75  $\mu\text{l}$  of 1 N NaOH. The quenched reaction mixture (160  $\mu\text{l}$ ) was transferred to 1 ml of 5% trichloroacetic acid and boiled to remove unincorporated <sup>14</sup>CO<sub>2</sub>. After cooling, 5 ml of Ecolite (PerkinElmer Life Sciences) was added, and the <sup>14</sup>CO<sub>2</sub> incorporated was determined in a Beckman LS 9800 counter.

#### RESULTS

**Characterization of a Putative *Drosophila*  $\gamma$ -Carboxylase mRNA Sequence**—Poly(A)<sup>+</sup> RNA from adult *Drosophila* was used as template for the reverse transcription of cDNA using an oligo(dT) primer. An initial segment of cDNA was amplified and sequenced using  $\gamma$ -carboxylase primers encoding amino acid sequences highly conserved between *Drosophila* sequences from the Berkeley *Drosophila* Genome Project (BDGP) and all mammalian enzymes; the 5'-oligonucleotide primer (primer 1; Table I) corresponded to amino acids 395–402, YGYSWDDMM, and the 3'-primer (primer 2; Table I) corresponded to amino acids 465–471, IYFDIWC (the amino acid positions correspond to human  $\gamma$ -glutamyl carboxylase sequence). The PCR product was cloned, and the nucleic acid sequence was determined; the

<sup>1</sup> The abbreviations used are: PCR, polymerase chain reaction; GC,  $\gamma$ -carboxylase; BDGP, Berkeley *Drosophila* Genome Project;  $\gamma$ -CRS,  $\gamma$ -carboxylation recognition signal sequence; GFP, green fluorescent protein; S2, Schneider 2.

TABLE I  
Primers used in PCR to identify *Drosophila*  $\gamma$ -glutamyl carboxylase

The primer pairs used in PCR amplification (see Fig. 2) were 1 and 2, 1 and Q<sub>0</sub>, 3 and Q<sub>0</sub>, 5 and 4, 5 and 7, 6 and 7. RACE, rapid amplification of cDNA ends.

Primer	Nucleotides	Sequence
1	1361–1384	TAYGGITAYTCATGGGAYATGATG
2	1599–1581	CACCAGATGTCGAAGTAGA
3	1513–1532	GCAGTAYGCCAGGTGCATCG
4	1532–1513	CGATGCACCTGGCCTACTGC
5	567–587	GTGGAATAACCACAGCTATCT
6	2–22	TGCAAAGGACGTGTTCTTTTC
7	2510–2419	ATCAGTTTATCACCATAC
8	Q <sub>T</sub> (3'-RACE primer)	CCAGTGAGCAGAGTGACGAGGACTCGAGCTCAAGCTTTTTTTTTTTTTTTT
9	Q <sub>0</sub>	CCAGTGAGCAGAGTGACG

sequence obtained was identical to a DNA sequence in the *Drosophila* genome (BDGP, accession number AC005557).

We carried out further PCR analysis of the cDNA using the primers indicated schematically in Table I and Fig. 2A. The 3'-end of the transcript to the poly(A) addition site was determined using 3'-rapid amplification of cDNA ends. Polyadenylation takes place 20 bases downstream of a consensus Poly(A) signal, AATAAA. The remainder of the cDNA was characterized by PCR amplification using primers shown in Table I; PCR products were cloned, and overlapping sequences were combined to yield the cDNA shown schematically in Fig. 2A.

The cDNA sequence encodes an open reading frame of 670 amino acids. Examination of the genomic sequence revealed two notable differences from the cDNA. First, there are two short introns. A schematic comparison of the genomic and cDNA, shown in Fig. 2B, illustrates the position of these introns. The nucleic acid sequences at the splice junction are shown in Fig. 2C. Second, an as yet uncharacterized processing event removes 6 nucleotides from the cDNA (Fig. 2D).

The locations of the two introns are conserved between *Drosophila* and mammals. As is generally found when comparing *Drosophila* with mammalian introns (30), the *Drosophila* introns are significantly shorter (for intron I, 58 versus 2204 nucleotides; for intron II, 72 versus 646 nucleotides). Comparison of the amino acid sequences flanking *Drosophila* intron II and human intron VII is shown in Fig. 2E. There are a total of 14 introns in both the rat (26) and the human  $\gamma$ -carboxylase genes (31); therefore, *Drosophila* has both fewer and shorter introns compared with the mammalian gene. Fig. 3A shows a schematic of amino acid homology between human and *Drosophila*  $\gamma$ -carboxylase.

The surprising finding from the sequencing described above is that an uncharacterized mechanism of RNA processing results in the deletion of 6 nucleotides that would have been present if the DNA were faithfully transcribed. The deletion does not change the amino acid homology to the mammalian enzyme at this site. This region of cDNA was sequenced at least three times as parts of PCR amplification products synthesized by different primer pairs. To confirm the genomic sequence of our strain, we directly sequenced the genomic DNA from the *Drosophila* strain that was the source of our cDNA. It was identical to that from BDGP, which confirmed that the deletion of 6 nucleotides was real and not an artifact of strain differences or cloning or sequencing errors.

With regard to the protein length, the alignment in Fig. 3B shows that although the *Drosophila* enzyme has 17 additional amino acids at the N terminus compared with the mammalian enzymes, it is significantly shorter at the C-terminal end. All of the mammalian enzymes are longer (758 versus 670 amino acids). A recent deletion analysis (32) of the bovine enzyme suggests that small deletions at the C terminus may be tolerated by the wild-type mammalian enzyme. Interestingly, a




FIG. 2. A, schematic of PCR products. Coordinates refer to the genomic sequence. Positions of initiation and termination codons, introns, processing site (*DELETED*), and Poly(A) addition signals are shown. B, schematic comparison of genomic and cDNA maps of *Drosophila*  $\gamma$ -carboxylase. C, nucleic acid and amino acid sequences at the splice junctions of *Drosophila* introns I and II. D, cDNA sequences at the processing site (*DELETED*). E, comparison of amino acid sequences flanking *Drosophila* intron II and human intron VII. Amino acids at the splice junction are shown in bold.

deletion that resulted in a bovine enzyme that was 676 amino acids in length had lower enzymatic activity (15-fold lower with respect to  $\gamma$ -carboxylation and 400-fold lower than for vitamin K epoxidation). It remains to be determined whether other *Drosophila* subunits are necessary to compensate for the shorter length of the *Drosophila* open reading frame.

**Northern Blot Analysis**—Fig. 4 shows the results of Northern blot analysis. *Drosophila*  $\gamma$ -glutamyl carboxylase mRNA is ~2.7 kb in size (Fig. 4A) and is predominantly expressed in 12–24-h embryos (Fig. 4B). (However, the more sensitive *in situ* hybridization experiments presented below reveal the presence of carboxylase mRNA throughout embryogenesis.) Ribosomal protein rp49 mRNA was also monitored in these experiments






FIG. 4. Northern blot analysis. A, total RNA from *Drosophila* embryos (0–24 h) probed with  $^{32}\text{P}$ -labeled antisense  $\gamma$ -carboxylase RNA. Molecular weight standards represent ethidium bromide-stained RNA Millennium markers (Ambion). B and C, analysis of developmental expression of  $\gamma$ -carboxylase (B) and rp49 (C) RNAs. Lanes containing RNA from embryos at 0–2, 2–4, 4–8, 12–16, and 16–24 h of development, larval stages 1–3, pupae, and adult flies are indicated (the lane between 4–8-h and 12–16-h embryo samples is blank).




FIG. 5. Ubiquitous transcription of  $\gamma$ -carboxylase in *Drosophila* embryos. *In situ* hybridization to stage 5 (A and B) and 16 (C and D) whole-mount embryos with digoxigenin labeled strand-specific probes is shown. Ubiquitous gene expression is visible when the antisense strand is used as probe (B and D) but not when the sense strand is used (A and C).

TABLE II  
Carboxylase activity in transfected *Drosophila* S2 cells

Enzyme <sup>a</sup>	$^{14}\text{CO}_2$ incorporated
	pmol
pRmHa-3.GC*-transfected S2 cell lysate	
Complete reaction	83
Minus vitamin K	4
Minus cell lysate	4
Mock-transfected S2 cell lysate	
Complete reaction	4

<sup>a</sup> Cell lysates from  $2 \times 10^5$  cells were used in each reaction.

served motif in this region has been suggested by Begley *et al.* (37). However, nucleic acid sequences reported here, by Li *et al.* (20) and by the BDGP (AC005557), are not consistent with the suggestion. The sequences reported here, by Li *et al.* (20) and by the BDGP (AC005557), are  $^{385}\text{GYNNWTNGLYGYSWDMMVH}^{404}\text{SYDTLQTSIQVVD} \dots$ , whereas the sequences at this site reported by Begley *et al.* (37) are  $^{385}\text{GYNNWTNGLYGYSWDMMVH}^{404}\text{SRSHQHVKITRYD}$ .

A recently elucidated hereditary disease further emphasizes the functional significance of the total conservation of amino acid sequence in this region: mutation of residue 395 (Leu→Arg) in the human enzyme (38) results in a clinical syndrome characterized by a general deficiency of blood clotting. Interestingly, this clinical condition can be treated satisfactorily by an infusion of high doses of vitamin K, consistent with an enzymatic defect in the affinity of the enzyme for its substrate, reduced vitamin K. Thus, the comparison between *Drosophila* and mammalian enzymes may have helped define a conserved site involved in the binding of reduced vitamin K, consistent with more conventional biochemical studies.

Heterologous  $\gamma$ -CRS sequences are not or are poorly recognized by the  $\gamma$ -glutamyl carboxylases.  $\gamma$ -CRS containing *Conus* substrate, proconantokin G, is poorly carboxylated by the bovine enzyme, whereas a peptide, factor IX-18–41, which consists of the propeptide and all normally carboxylated residues of the vitamin K-dependent clotting protein factor IX, is not carboxylated by the *Conus* enzyme (18). The poor carboxylation of -20Y by the *Drosophila* enzyme further strengthens the suggestion that the enzymes have evolved to recognize their cognate  $\gamma$ -CRSs. This is also supported by the observation of Li *et al.* (20), who found that the propeptide of human blood coagulation factor IX did not stimulate carboxylation by the *Drosophila* enzyme. Because the *Drosophila* and human  $\gamma$ -glutamyl carboxylases share considerable sequence homology, it should be possible to identify substrate binding domains by studying carboxylation using chimeric enzymes.

Because  $\gamma$ -carboxylated molecules may serve as signals for growth and differentiation, differential regulation of  $\gamma$ -carboxylation may operate at multiple levels during development. Control may be at the level of synthesis of  $\gamma$ -carboxylase mRNA or its translation, or both. Although mRNA may be present, enzyme activity may not be obvious. Future experiments will be aimed at determining possible differences among levels of mRNA, expressed protein, and activity by immunological methods (for protein) and  $\gamma$ -carboxylase assay (for activity).

$\gamma$ -Carboxyglutamate-containing proteins isolated to date are extracellular proteins.  $\gamma$ -Carboxyglutamate interacts with  $\text{Ca}^{2+}$ , induces a conformational change in the protein, and facilitates binding to membrane phospholipids. A number of  $\gamma$ -carboxyglutamate-containing vitamin K-dependent proteins (thrombin, factor Xa, protein S, and Gas6) are ligands for cell surface receptors. Interaction with the receptors induces cellular proliferative responses (39, 40). In *Drosophila*, high levels of  $\gamma$ -carboxylase RNA are detected in late stage embryos. During this period, a variety of developmental and morphogenetic events occur, among them cuticle deposition and central nervous system, peripheral nervous system, and gut differentiation. It is conceivable that some of the gene products signaling these events are  $\gamma$ -carboxylated and serve as ligands for corresponding receptors. The effects of  $\gamma$ -carboxylase knockout in flies will enable a systematic study of probable targets for this post-translational modification.

**Acknowledgment**—We thank the DNA Sequencing Facility at the University of Utah. The Peptide Sequencing facility at the University of Utah is supported by NCI Grant 5p30CA42014.

#### REFERENCES

1. Stenflo, J., Fernlund, P., Egan, W., and Roepstorff, P. (1974) *Proc. Natl. Acad. Sci. U. S. A.* **71**, 2730–2733
2. Price, P. A., and Williamson, M. K. (1985) *J. Biol. Chem.* **260**, 14971–14975
3. Pan, L. C., and Price, P. A. (1985) *Proc. Natl. Acad. Sci. U. S. A.* **82**, 6109–6113
4. Manfioletti, G., Brancolini, C., Avanzi, G., and Schneider, C. (1993) *Mol. Cell. Biol.* **13**, 4976–4985
5. Kulman, J. D., Harris, J. E., Haldeman, B. A., and Davie, E. W. (1997) *Proc.*

- Natl. Acad. Sci. U. S. A.* **94**, 9058–9062
6. McIntosh, J. M., Olivera, B. M., Cruz, L. J., and Gray, W. R. (1984) *J. Biol. Chem.* **259**, 14343–14346
  7. Craig, A. G., Bandyopadhyay, P., and Olivera, B. M. (1999) *Eur. J. Biochem.* **264**, 271–275
  8. Hauschka, P. V., Mullen, E. A., Hintsch, G., and Jazwinski, S. (1988) in *Current Advances in Vitamin K Research* (Suttie, J. W., ed) pp. 237–243, Science Publishers, New York
  9. Blandl, T., Zajicek, J., Prorok, M., and Castellino, F. J. (1997) *Biochem. J.* **328**, 777–783
  10. Rigby, A. C., Baleja, J. D., Furie, B. C., and Furie, B. (1997) *Biochemistry* **36**, 6906–6914
  11. Warder, S. E., Chen, Z., Zhu, Y., Prorok, M., Castellino, F. J., and Ni, F. (1997) *FEBS Lett.* **411**, 19–26
  12. Skjaebaek, N., Nielsen, K. J., Lewis, R. J., Alewood, P., and Craik, D. J. (1997) *J. Biol. Chem.* **272**, 2291–2299
  13. Myers, R. A., McIntosh, J. M., Imperial, J., Williams, R. W., Oas, T., Haack, J. A., Hernandez, J.-F., Rivier, J., Cruz, L. J., and Olivera, B. M. (1990) *J. Toxicol. Toxin Rev.* **9**, 179–202
  14. Soriano-Garcia, M., Padmanabhan, K., de Vos, A. M., and Tulinsky, A. (1992) *Biochemistry* **31**, 2554–2566
  15. Freedman, S. J., Furie, B. C., Furie, B., and Baleja, J. (1995) *Biochemistry* **34**, 12126–12137
  16. Sunnerhagen, M., Forsen, S., Hoffren, A. M., Drakenberg, T., Teleman, O., and Stenflo, J. (1995) *Nat. Struct. Biol.* **2**, 504–509
  17. Mann, K. G., Nesheim, M. E., Church, W. R., Haley, P., and Krishnawamy, S. (1990) *Blood* **76**, 1–16
  18. Stanley, T. B., Stafford, D. W., Olivera, B. M., and Bandyopadhyay, P. K. (1997) *FEBS Lett.* **407**, 85–88
  19. Bandyopadhyay, P. K., Colledge, C. J., Walker, C. S., Zhou, L.-M., Hillyard, D. R., and Olivera, B. M. (1998) *J. Biol. Chem.* **273**, 5447–5450
  20. Li, T., Yang, C.-T., Jin, D., and Stafford, D. W. (2000) *J. Biol. Chem.* **275**, 18291–18296
  21. Chomczynski, P., and Sacchi, N. (1987) *Anal. Biochem.* **162**, 156–159
  22. Sambrook, J., Fritsch, E. F., and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, NY
  23. Frohman, M. A. (1994) in *The Polymerase Chain Reaction* (Mullis, K. B., Ferré, F., and Gibbs, R. A., eds) pp. 14–37, Birhauser, Boston
  24. Wu, S.-M., Cheung, W.-F., Frazier, D., and Stafford, D. W. (1991) *Science* **254**, 1634–1636
  25. Rehemtulla, A., Roth, D. A., Wasley, L. C., Kuliopulos, A., Walsh, C. T., Furie, B., Furie, B. C., and Kaufman, R. J. (1993) *Proc. Natl. Acad. Sci. U. S. A.* **90**, 4611–4615
  26. Romero, E. E., Velazquez-Estades, L. J., Deo, R., Schapiro, B., and Roth, D. A. (1998) *Exp. Cell Res.* **243**, 334–346
  27. Lehmann, R., and Tautz, D. (1994) in *Methods in Cell Biology*, (Goldstein, L. S. B., and Fyrberg, E. A., eds) Vol. 44, pp. 575–598, Academic Press, New York
  28. Bunch, T. A., Grinblat, Y., and Goldstein, L. S. B. (1988) *Nucleic Acids Res.* **16**, 1043–1061
  29. Esmon, C. T., Sadowski, J. A., and Suttie, J. W. (1975) *J. Biol. Chem.* **250**, 4744–4748
  30. Deutsch, M., and Long, M. (1999) *Nucleic Acids Res.* **27**, 3219–3228
  31. Stafford, D. W., Frazier, L. D., Fu, Y. Y., High, K. A., Chu, K., Sanchez-Vega, B., and Solera, J. (1997) *Blood* **89**, 4058–4062
  32. Roth, D. A., Whirl, M. L., Velazquez-Estades, L. J., Walsh, C. T., Furie, B., and Furie, B. C. (1995) *J. Biol. Chem.* **270**, 5305–5311
  33. Andres, A. J., and Cherbas, P. (1992) *Development* **116**, 865–876
  34. Stronach, B. E., Siegrist, S. E., and Beckerle, M. C. (1996) *J. Cell Biol.* **134**, 1179–1195
  35. Kuliopulos, A., Cieurzo, C. E., Furie, B., Furie, B. C., and Walsh, C. T. (1992) *Biochemistry* **31**, 9436–9444
  36. Bush, K. A., Stenflo, J., Roth, D. A., Czerwicz, E., Harrist, A., Begley, G. S., Furie, B. C., and Furie, B. (1999) *Biochemistry* **38**, 14660–14666
  37. Begley, G. S., Furie, B. C., Czerwicz, E., Taylor, K. L., Furie, G. L., Bronstein, L., Stenflo, J., and Furie, B. (2000) *J. Biol. Chem.* **275**, 36245–36249
  38. Brenner, B., Sanchez-Vega, B., Wu, S., Lanir, N., Stafford, D. W., and Solera, J. (1998) *Blood* **92**, 4554–4559
  39. Esmon, C. T. (1995) *Curr. Biol.* **5**, 743–746
  40. Crosier, K. E., and Crosier, P. S. (1997) *Pathology* **29**, 131–135