

Priorities for hydrocephalus research: report from a National Institutes of Health–sponsored workshop

MICHAEL A. WILLIAMS, M.D.,^{1,2} JAMES P. MCALLISTER, PH.D.,^{3,4} MARION L. WALKER, M.D.,⁴ DORY A. KRANZ, M.A.,⁵ MARVIN BERGSNEIDER, M.D.,⁶ MARC R. DEL BIGIO, M.D., PH.D.,⁷ LAUREL FLEMING, B.A.,⁸ DAVID M. FRIM, M.D.,⁹ KATRINA GWINN, M.D.,¹⁰ JOHN R. W. KESTLE, M.D.,⁴ MARK G. LUCIANO, M.D., PH.D.,¹¹ JOSEPH R. MADSEN, M.D.,⁸ MARY LOU OSTER-GRANITE, PH.D.,¹² AND GIOVANNA SPINELLA, M.D.¹³

¹Department of Neurology, Johns Hopkins Medical Institutions, Baltimore; ²LifeBridge Health Brain and Spine Institute, Adult Hydrocephalus Center, Baltimore, Maryland; ³Department of Neurosurgery, Wayne State University, Detroit, Michigan; ⁴Department of Neurosurgery, University of Utah, Salt Lake City, Utah; ⁵Hydrocephalus Association, San Francisco; ⁶Department of Neurosurgery, University of California, Los Angeles, California; ⁷Department of Pathology, University of Manitoba, Winnipeg, Canada; ⁸Department of Neurosurgery, Harvard University School of Medicine, Boston, Massachusetts; ⁹Department of Neurosurgery, University of Chicago, Illinois; ¹⁰National Institute of Neurological Disorders and Stroke, Bethesda, Maryland; ¹¹Department of Neurosurgery, Cleveland Clinic, Cleveland, Ohio; ¹²National Institute of Child Health and Human Development; and ¹³National Institutes of Health, Bethesda, Maryland

Object. Treatment for hydrocephalus has not advanced appreciably since the advent of cerebrospinal fluid (CSF) shunts more than 50 years ago. Many questions remain that clinical and basic research could address, which in turn could improve therapeutic options. To clarify the main issues facing hydrocephalus research and to identify critical advances necessary to improve outcomes for patients with hydrocephalus, the National Institutes of Health (NIH) sponsored a workshop titled “Hydrocephalus: Myths, New Facts, and Clear Directions.” The purpose of this paper is to report on the recommendations that resulted from that workshop.

Methods. The workshop convened from September 29 to October 1, 2005, in Bethesda, Maryland. Among the 150 attendees was an international group of participants, including experts in pediatric and adult hydrocephalus as well as scientists working in related fields, neurosurgeons, laboratory-based neuroscientists, neurologists, patient advocates, individuals with hydrocephalus, parents, and NIH program and intramural staff. Plenary and breakout sessions covered injury and recovery mechanisms, modeling, biomechanics, diagnosis, current treatment and outcomes, complications, quality of life, future treatments, medical devices, development of research networks and information sharing, and education and career development.

Results. The conclusions were as follows: 1) current methods of diagnosis, treatment, and outcomes monitoring need improvement; 2) frequent complications, poor rate of shunt survival, and poor quality of life for patients lead to unsatisfactory outcomes; 3) investigators and caregivers need additional methods to monitor neurocognitive function and control of CSF variables such as pressure, flow, or pulsatility; 4) research warrants novel interdisciplinary approaches; 5) understanding of the pathophysiological and recovery mechanisms of neuronal function in hydrocephalus is poor, warranting further investigation; and 6) both basic and clinical aspects warrant expanded and innovative training programs.

Conclusions. The research priorities of this workshop provide critical guidance for future research in hydrocephalus, which should result in advances in knowledge, and ultimately in the treatment for this important disorder and improved outcomes in patients of all ages. (DOI: 10.3171/PED-07/11/345)

KEY WORDS • adult • basic biomedical research • children • clinical research • diagnostic criteria • hydrocephalus • infant • National Institutes of Health • neurosurgical education • normal pressure hydrocephalus • pediatric neurosurgery • therapeutic intervention

Abbreviations used in this paper: CBF = cerebral blood flow; CSF = cerebrospinal fluid; CT = computed tomography; ETV = endoscopic third ventriculostomy; ICP = intracranial pressure; MR = magnetic resonance; NIH = National Institutes of Health; NINDS = National Institute of Neurological Disorders and Stroke; NPH = normal-pressure hydrocephalus; R_{out} = CSF outflow resistance.

HYDROCEPHALUS is the abnormal primary enlargement of the cerebral ventricles that results from impaired CSF secretion, circulation, or resorption. It is distinct from “hydrocephalus ex vacuo,” the abnormal secondary enlargement of the cerebral ventricles resulting from loss of cerebral tissue (for example, due to cerebral atrophy).

Most commonly recognized as a disorder of infants and children,^{73,104} hydrocephalus also affects adolescents, young and middle-aged adults,^{31,71,139} and the elderly.^{82,118,121,126,152}

The first effective treatment for hydrocephalus, the shunt, was developed 50 years ago by John Holter, in collaboration with the neurosurgeon Eugene Spitz.⁵ The shunt revolutionized the care of patients with hydrocephalus; shunt surgery is now the most common procedure pediatric neurosurgeons perform. With this dramatic success, the search for other approaches to treatment slowed.

Despite improvements in neuroimaging, shunt valve technology, and ETV, little has changed in the diagnostic and treatment approaches toward hydrocephalus in the last 50 years. Surprisingly little clinical or basic research has focused on hydrocephalus, aside from studies to test the effectiveness of new shunts (often to achieve US Food and Drug Administration approval). Frequent complications and associated morbidity often make treatment outcomes unsatisfactory.⁸⁰ The infrastructure for clinical care in the US is largely limited to pediatric neurosurgeons, and fewer than 10 centers have expertise for diagnosing and treating adult hydrocephalus. No standard mechanism exists for the smooth transition of care for individuals with hydrocephalus from pediatric to adult specialists.

Hydrocephalus is far from cured, and there is a pressing need for basic and clinical research to improve our understanding of this complex disorder. Hydrocephalus is sometimes misconstrued as a simple disorder of CSF circulation—a “plumbing problem”—when it is more properly viewed as a brain disorder. Because hydrocephalus signs and symptoms can resolve in days or weeks after existing for months or years, a novel form of reversible neuronal and glial injury may be involved. At present, the basic injury and recovery mechanisms are not adequately understood.

In recognition of the gaps in our knowledge, and the benefits that could emerge from focused research, the NIH convened a workshop titled “Hydrocephalus: Myths, New Facts, and Clear Directions” in Bethesda, Maryland, from September 29 to October 1, 2005. More than 150 neurosurgeons, neurologists, laboratory-based scientists, patient advocates, patients, parents, and program staff from five institutes and one office within the NIH assembled for a day and a half of plenary discussions that summarized the current knowledge, challenged existing dogma and mythology, and identified critical gaps in research and clinical treatment. Participants divided into three breakout groups to answer key questions (Appendices 1 and 2) with the goal of developing research priorities and collaborative opportunities.

The priorities for research (Table 1) resulting from the 2005 NIH Hydrocephalus Workshop should set in motion a broad, interdependent array of research in hydrocephalus that we hope will yield important discoveries fundamental to our understanding of brain function and CSF physiology and ultimately lead to therapeutic intervention strategies and improved outcomes for patients of all ages.

Clinical, Research, and Education Infrastructure

The research priorities begin with recommendations for research infrastructure (Table 1) that all three breakout groups identified independently as a need for all types of research (clinical, basic, translational, and interdisciplinary),

TABLE 1

Research priorities for hydrocephalus

<i>Hydrocephalus Research Infrastructure</i>	
Creation of research networks	Basic, clinical, translational, & interdisciplinary research
Facilitation of information sharing	Specimen banks: brain, CSF, genetic material, shunts, images, physiological measurements
	Registries: clinical trials, epidemiological data, longitudinal studies
Education & career development for researchers	Scientist & clinician/scientist education
<i>Basic Research</i>	
Injury & recovery mechanisms in human volunteers or animal models	Molecular, cellular, genetic, & systems physiology
	Role of stem cells & neural progenitor cells
	Role of inflammation, tissue matrix, blood–brain barrier, biomarkers
Anatomy & physiology of CSF secretion, circulation, resorption, function	
Development of appropriate animal models for age, acuity of onset, & mode of induction factors, including congenital & acquired hydrocephalus	Ex vivo models (brain slices, tissue culture)
	Transgenic models
Development & validation of theoretical & biomechanical modeling	CBF/CSF pulsatility, compartment models, ICP gradients, & monitoring
	Physiological effects of CSF diversion, including slit-ventricle syndrome
<i>Clinical Research</i>	
Diagnosis	Establishment & validation of diagnostic criteria at all stages of life
	Improvement & validation of imaging & noninvasive testing techniques
Current treatment, outcomes, complications	Epidemiology & population impact
Surgical treatment	Establishment & validation of treatment criteria at all stages of life
	Prevention of infection & device or treatment failure
	Effects of device design, biomaterials, & technique on outcomes
Clinical evaluation of brain/neuronal dysfunction & recovery	Development & validation of clinical conventions or templates for assessing neurological & behavioral outcomes (for example, cognitive, gait, motor, continence, activities of daily living)
	Correlation of clinical outcomes w/ imaging & noninvasive techniques
	Effects of infection & device or treatment failure on outcomes
Health services research	Education of patients, public, physician communities
	Determination of extent of need for centers of excellence
	Determination of extent of need for longitudinal care for children & adults w/ chronic hydrocephalus
	Development of age-appropriate quality of life instruments
<i>Future Treatments</i>	
	Development of adjunctive (pharmacological) therapies
	Development of novel medical devices & biomaterials

which should facilitate the creation of research networks, data sharing, and education for future researchers.

Virtually no organized clinical, research, or educational infrastructure for hydrocephalus currently exists. Nevertheless, there is a need, given the fact that the cost of treating hydrocephalus is estimated at 1 billion dollars annually,¹⁴⁴ and there is evidence that treating adults with hydrocephalus who are older than 65 years of age can lower national 5-year Medicare expenditures by as much as \$184.3 million.¹⁸⁴

Priorities for hydrocephalus research

Creation of a Clinical Research Network for Hydrocephalus

Between 1996 and 2006, there were only eight publications about multicenter hydrocephalus trials (according to a search of Medline [accessed September 2006]). There is a need to establish clinical research networks such as the Children's Oncology Group, which is funded by the National Cancer Institute. Such multicenter involvement encourages pooling of data, tissue specimens, and knowledge as well as potential treatment innovation and research of clinical outcomes. Most data on the longevity of CSF shunts, for example, reside with shunt manufacturing companies. The ability to compare different shunt models across a large population would be beneficial. A multiinstitutional clinical research network could help answer basic questions, such as the outcome of treated compared with untreated hydrocephalus, or facilitate comparisons of surgical strategies and outcomes, such as shunts with or without antisiphoning devices or shunts compared with ETV. Investigators could use the research network database for the generation of hypotheses to be used in more focused research.

Creation of a Clinical Care and Research Training Infrastructure

There is currently no formal postgraduate clinical or research training program in the care of hydrocephalus. Much of the training resides within pediatric neurosurgery fellowships, not all of which have a clinical or research focus on hydrocephalus. This ad hoc system generally does not encompass adult hydrocephalus, nor does it include other specialists, such as neurologists and clinically oriented neuroscientists. Thus, the goal of reaching a "critical mass" of interdisciplinary clinical and research scientists studying hydrocephalus is a major challenge. It is plausible that the lack of critical mass or individual research mentors is a disincentive for young investigators to enter a career in hydrocephalus research.

Creation of Infrastructure to Train Basic Science Researchers

It is arguable that in the field of hydrocephalus research there may be even fewer basic scientists than there are clinicians or clinical scientists. Considering that the pathophysiological processes in this disease are diverse and poorly understood, this field is an unparalleled opportunity for clinical or basic researchers. There should be incentives for such clinicians and scientists to concentrate their career paths toward hydrocephalus research. The establishment of funded postdoctoral fellowships could provide an incentive to obtain training, as could targeted faculty positions or endowed chairs that would provide stable and productive environments for junior researchers.

Hydrocephalus Patient Advocacy

Some diseases and disorders have garnered focused federal funding for research due to the efforts of advocacy groups composed of philanthropic leaders, patients, and their families. The Hydrocephalus Association and other volunteer organizations with an interest in hydrocephalus that played an important role in the NIH workshop have filled that role in many ways. For example, the Hydrocephalus

Association has made a deliberate choice to promote federal funding for hydrocephalus research rather than fund research on its own as similar organizations have done (for example, the Multiple Sclerosis Society and the Amyotrophic Lateral Sclerosis Society). Patient advocacy groups also provide information to patients and their families with regard to diagnostic and treatment options and ongoing clinical research trials. Collaboration with patient advocacy groups should be encouraged, and the emerging hydrocephalus research community should view such cooperation as a highly valuable component of any complete clinical and research program.

Priorities for Basic Research

Priorities for basic research (Table 1) focus on the injury and recovery mechanisms of hydrocephalus, ranging from the genetic and molecular levels to animal models and systems physiology approaches. Theoretical modeling of the effects of CSF, CBF, and ICP on normal and abnormal physiology is also needed.

Injury and Recovery Mechanisms

The signs and symptoms of hydrocephalus, particularly in adults (but also in children), may be present for months or years, and yet resolve in days or weeks. Consequently, they represent a unique and clinically important type of neuronal injury and recovery. There is inadequate understanding of the mechanisms that lead to the initiation, progression, and persistence of neuronal, glial, and vascular dysfunction in hydrocephalus, as well as the mechanisms by which treatment reverses this injury. The application of contemporary methods and techniques in basic neurobiology will provide critical insights into both the injury and recovery mechanisms of hydrocephalus, and we hope that this will lead to improved diagnostic, prognostic, and treatment approaches.

Hydrocephalus has many known causes, including congenital malformations, hemorrhage, infection (primarily meningitis), and trauma, among others. However, most of the available knowledge pertains to secondary injuries and responses,^{41,48,123} including neuronal and glial cell death, axonal and dendritic degeneration, astroglial and microglial activation and proliferation, neurochemical alterations, decreases in CBF, and impaired cellular metabolism.^{41,44-46,48,92} Animal models of hydrocephalus include natural (congenital) forms in rodents or forms induced in many species by injection of materials such as blood, kaolin, silicone, or acrylic polymers that obstruct CSF flow either through an intense subarachnoid inflammatory response or a mechanical barrier.^{32,88,140}

Regardless of the cause, dilation of the lateral ventricles results in several types of brain injury. The ependymal lining is stretched and eventually lost. Cells in the subependymal zone proliferate, giving rise to reactive astrocytes. Oligodendrocyte production of myelin is retarded, and these cells eventually die. Reactive astroglial and microglial changes occur in the periventricular white matter but also in the cortical and subcortical gray matter. The extracellular fluid compartment is enlarged in the white matter but compressed in the cortical laminae. Periventricular axons are stretched and can be destroyed.^{40,68,109,110} The prominent white matter damage suggests that hydrocephalus could be

considered a type of subcortical disconnection syndrome.⁵⁰ Disconnected neurons undergo atrophy of dendritic processes and synapses and may die.^{19,85,86,109,110,124} Decreases in neurotransmitters and neuromodulators are common,^{30,91,113,132} and reductions in energy metabolites occur within days of the onset of ventriculomegaly.^{49,87,99,132} Hydrocephalus adversely affects the cerebral vasculature, CBF, and metabolism,^{6,23,43,61,143,153,179,189} and there may be indirect effects on CBF via cardiac suppression.^{53,58}

A number of factors modify the pathogenesis of brain injury in hydrocephalus, including the age of the individual and the state of brain maturation at onset, the rate of ventricular dilation, duration of stable ventriculomegaly, cause of hydrocephalus, and coexistence of other diseases. In the developing brain, enlargement of the ventricles and stretching of the ependymal lining raises complex questions about reduction of germinal cell proliferation, alteration of synaptic connections, and brain development.^{106,133,134,142,175} There has been no systematic study of the effect of repeated episodes of ventricular dilation and contraction associated with shunt obstruction and revision. It is plausible that repeated injury to neurons and glia is cumulative, and the capacity to reverse such injury may be lost.¹²³ The calcium-mediated proteolytic changes in axons are similar to those in stroke and trauma.⁴⁰ In vitro neuron studies might be used to determine how changes in CSF and extracellular fluid composition are toxic to axons and neurons in hydrocephalus. Understanding the secondary role of microglial inflammation in white matter is also critical.^{116,174}

Gross restoration of the brain, blood, interstitial fluid, and CSF volumes and their volume relationships can occur quickly after CSF shunting;⁴³ however, reduction in ventricle size is not always necessary for clinical improvement, particularly in adults. Thus, some component of the quickly reversible brain dysfunction may result from intracellular neuronal or glial changes (for example, biochemical changes or gene induction). Within a span of hours to days, CSF shunting likely corrects white matter CBF and also may divert clearance of waste products to alternate absorption sites. Relief of metabolic and physical stresses on axons may prevent injury progression. In young hydrocephalic animals, early shunt placement is associated with a “catch-up” in myelin production⁴⁷ and the restoration of axoplasmic transport in cortical connections within days.⁶⁸

A subset of fetal or childhood-onset hydrocephalus is related to developmental anomalies that are sometimes attributed to genetic abnormalities. However, the role of genetic factors in the development of hydrocephalus overall is not well understood. The only hydrocephalus gene identified in humans, the X-linked *LI-NCAM* mutation, is associated with many brain anomalies.^{69,155,167,177} Considering that a significant proportion of cases of hydrocephalus in humans occurs as part of a more widespread developmental anomaly, many defective genes may act singly or (more likely) in combination.¹⁵⁹ Studies of gene expression in models of either induced or genetic hydrocephalus may identify molecular mechanisms of injury and recovery.^{4,131,136} A database that listed rodents with hydrocephalus, their genetic characterization, other phenotypic data, and associated publications would be an important part of the research infrastructure.

Many more hereditary forms of hydrocephalus occur in animals than in humans, including those found in the *H-Tx*

rat, the *LEWS/Jms* rat, and the *hy-3* mouse.^{8,24,25,95–98,100,101,122,151,154,178,188} Transforming growth factor- β_1 may be involved in posthemorrhagic or postmeningitic hydrocephalus,^{168,169,183} and upregulation of transforming growth factor- β_1 causes abnormal extracellular matrix in the CSF flow pathways.⁷² Inducing hydrocephalus in mutant animals that do not otherwise develop this disorder can be instructive. For example, mice lacking aquaporin 4 develop hydrocephalus more rapidly after kaolin injection,¹⁸ and rats with kaolin-induced hydrocephalus upregulate aquaporin 4.¹¹⁷ The relationship of inherited abnormalities of ciliary function to hydrocephalus is not understood.^{3,155,177}

There are no models with which to study the effects of chronic hydrocephalus or hydrocephalus arising across the age spectrum from brain development to senescence. Such models might provide a basis for treating the disorder in utero, or might lead to strategies to inhibit cerebral injury until treatment could be offered after birth.^{57,74,130} Similarly, understanding the effects of chronic ventriculomegaly on the brain may guide decisions to treat or not treat adults with “stable” hydrocephalus.¹⁸⁵ Models using senescent animals might improve understanding of the interaction of hydrocephalus with hypertensive microvascular disease and Alzheimer disease, which are common in elderly patients with idiopathic NPH.^{93,94,166,185}

In summary, despite progress in their identification over the past 30 years, the most important injury and recovery mechanisms in hydrocephalus remain elusive. This undoubtedly impedes the translational research necessary to advance the clinical management of hydrocephalus at all stages and ages of life. Research conducted using existing animal models should be expanded, and there is a need for new, refined, and more physiologically appropriate animal models.

Mathematical Modeling and Biomechanics

The study of the movement of CSF and interstitial fluid within the craniospinal compartment is fundamental to the understanding of hydrocephalus, and mathematical modeling has played a major role in its investigation.¹⁷¹ Two tenets have guided the modeling of CSF dynamics: the Monro-Kellie doctrine and CSF bulk flow.

The Monro-Kellie doctrine^{102,135} depicts the cranium as a rigid chamber with a fixed volume that contains three liquid (and therefore incompressible) components: brain, blood, and CSF. If the volume of one of the components increases, then there must be compensation by a decrease in the volume of the other components. These principles form the basis for a variety of intracranial parameters, such as compliance, elastance, and the pressure-volume index.^{82,83}

The concept of bulk flow of CSF has dominated the modeling of CSF circulation and the management of CSF-related disorders.^{1,33–37,62,64–67,76,107,108,127–129,137,143,157,158,173,176} since 1919, when Dandy³⁸ demonstrated the production of CSF by the choroid plexus. The rate of CSF production is thought to be constant, and CSF flows through the ventricles, exits into the subarachnoid space, and is absorbed through the arachnoid granulations at the superior sagittal sinus or the spinal rootlets. Absorption of CSF is thought to be linearly related to the pressure gradient between the subarachnoid space and the intraluminal space of the dural venous sinuses. Consequently, hydrocephalus is often modeled as a disturbance of CSF absorption that results in a

Priorities for hydrocephalus research

“back-up” of fluid that increases CSF pressure and enlarges the ventricles.

Although there is little disagreement that bulk CSF flow circulation exists, its characteristics and importance are in question. For example, the net bulk flow of CSF over the cerebral convexities cannot be demonstrated by high-resolution cine MR imaging.^{78,79} Cerebrospinal fluid may freely cross parenchymal capillaries and plexuses.^{89,156} The concentration of radioisotope along the superior sagittal sinus and the sacral subarachnoid space may reflect a dilution effect, rather than the movement of CSF carrying the radioisotope.⁷⁹ If increased R_{out} at the arachnoid granulations causes communicating hydrocephalus, there should be an increased pressure gradient between the subarachnoid space and the sagittal sinus lumen, but currently there is no evidence of this.^{61,165} In obstructive hydrocephalus in adults, there is no difference in R_{out} measured in the ventricles and the subarachnoid space.¹⁷³

Pulsatility Biomechanics. The bulk flow model does not account for the dynamic, pulsatile movement of CSF, although Hakim⁸¹ acknowledged its contribution. The conventional view of hydrocephalus as an imbalance of CSF production and absorption may be an oversimplification.⁷⁷⁻⁷⁹ Newer modeling approaches consider the intracranial compartment to be highly dynamic and complex, relating the movement of CSF to CBF hemodynamics.

The cerebral arteries and choroid plexus generate intracranial pulsatility.^{14-16,28,29,51,52} In fact, ventriculomegaly can be induced simply by inserting a balloon in the lateral ventricle that pulsates synchronously with arterial pulsations and increases CSF pulse pressure without changing CSF mean pressure.^{51,52} Whereas obstruction of CSF flow can cause acute hydrocephalus, chronic hydrocephalus may result from increased capillary pulse pressure that causes ventricular enlargement.

The hemodynamic theory of ICP physiology posits that the dynamic movement of CSF across the foramen magnum into the distensible spinal subarachnoid space is essential for pulsatile blood flow in the intracranial compartment,¹¹⁻¹³ and this has been demonstrated experimentally in the venous system.⁹⁰ It has been proposed that disruption of pulsatile CSF flow should perturb CBF dynamics, leading to CBF autoregulatory and other compensatory mechanisms that result in disorders of CSF circulation.^{59-61,115,179,180}

More recently, results of modeling support the suggestion that the intracranial contents oscillate at an exact “tuned” frequency, rather than passively responding to the pulsatile “hammer” of the incoming systolic blood pressure and flow. The oscillation of the intracranial contents may facilitate the “smooth” flow of blood through the intracranial compartment. There is preliminary evidence for one prediction based on the model, which is that a tuned oscillator should act as a filter, damping some flow and pressure waveform frequencies but not others.¹¹⁵

Finite Element Analysis Modeling. Finite element analysis has been used to identify brain regions that are particularly susceptible to strain and compression during progressive ventriculomegaly,¹⁴⁵ and results of that analysis support the suggestion that a relative reduction in intraparenchymal fluid pressure coupled with low tissue elasticity causes both ventricular enlargement and mechanical stress in the cortical mantle.¹⁴⁶

Further research of mathematical modeling and biome-

chanical approaches may yield important new insights into the pathophysiological mechanisms of hydrocephalus.

Priorities for Clinical Research

Clinical research priorities (Table 1) include the improvement of diagnostic tests and criteria and of treatment criteria for patients at all stages of life; improvement of surgical therapy to prevent infections and treatment failure, and understanding of the impact of these adverse events on brain function and clinical outcomes; and development of clinical rating methods and correlation with other noninvasive techniques for diagnosis and outcomes research.

Diagnostic Tests

Diagnostic Criteria for Children and Adults. The goal of diagnostic and prognostic testing is to determine the need for treatment, the nature of the treatment, outcomes, and the need for treatment modification. Current diagnostic tools include the patient history and physical examination, ultrasonography in infants, CT scanning and MR imaging of the brain, and tests of CSF circulatory physiology (infusion testing for R_{out} , ICP monitoring, or symptomatic response to CSF removal).

The diagnosis of hydrocephalus is obvious when the ventricles expand on serial images and these findings are accompanied by symptoms of increased ICP. Nevertheless, the diagnosis, classification, and decision to treat are not always straightforward; for example, differentiating between progressive hydrocephalus and “compensated” hydrocephalus in apparently asymptomatic children, or distinguishing hydrocephalus from the ventriculomegaly of cerebral atrophy (ex vacuo) in the elderly.

In spite of earlier suggestions that cortical mantle thickness predicted outcome,¹⁸⁶ ventricular size alone cannot guide the diagnosis or the decision to treat hydrocephalus.⁸⁴ Young children with enlarged ventricles may develop normal or even superior function. In the elderly, the fact that either cerebral atrophy or normal aging can result in ventricular enlargement makes it difficult to establish reliable standards for ventriculomegaly. Even the use of CSF pressure measurement as a diagnostic criterion is limited because adult or pediatric chronic hydrocephalus may not be accompanied by an elevation in pressure.

If hydrocephalus results from a disturbance of CSF absorption, then measurement of R_{out} by using infusion methods should be a sound basis for diagnosis, as has been proposed for idiopathic NPH.^{118,121,127} However, the usefulness of R_{out} measurement, especially in view of its technical difficulty and variability in techniques, is controversial as well.

The conundrum of diagnosis without a full understanding of pathophysiological mechanisms is demonstrated by decades of clinical research in which the “gold standard” for diagnosis is the response to surgical treatment itself. Guidelines for diagnosis and treatment of idiopathic NPH in the elderly have been established,¹²⁰ but there are still no guidelines for newborns, children, adolescents, and young or middle-aged adults.

Because disturbed CSF circulation contributes to hydrocephalus, detection of CSF flow abnormalities should be useful in diagnosis and treatment evaluation. Cine CSF flow MR imaging has been used to differentiate communicating

hydrocephalus (hyperdynamic aqueductal flow) from aqueductal stenosis (absent aqueductal flow).^{7,179} Flow studies can show the patency of the third ventricle fenestration after ETV. Cine CSF flow MR imaging is limited by its ability to quantify flow only in specific directions, fields, and velocities.

Hydrocephalus is a chronic disorder. Thus, the assessment of treatment failure is as important as the initial diagnosis. The diagnosis of shunt or ETV failure and recurrence of hydrocephalus can be straightforward, with demonstration of expanding ventricles and symptoms of increasing ICP. The challenge is to develop diagnostic procedures that are sensitive enough to predict treatment failure before it happens or that allow intervention to be offered before permanent injury occurs. Severe morbidity and death are still seen as the result of sudden treatment failure.

Because the symptoms of shunt failure may resemble common disorders such as a viral syndrome, emergency department visits, CT or MR imaging studies, or shunt taps are often ordered to avoid harm due to "failure to diagnose." Finally, shunts may be "functioning" but still cause chronic unphysiological overdrainage of CSF. Many of these patients have headaches, nausea, and other ongoing symptoms, and yet are considered successfully treated because the ventricles are not enlarged. Improved and more sensitive diagnostic methods are needed to detect suboptimal treatment and treatment failure.

Imaging and Noninvasive Testing. Beginning with pneumoencephalography,³⁸ neuroimaging has been important for the diagnosis and management of hydrocephalus. Ventriculomegaly remains the sine qua non of hydrocephalus, and CT and MR imaging offer superb anatomical detail. Transependymal flow on CT scanning or MR imaging suggests the presence of increased ICP. Large ventricles without transependymal flow in the presence of normal examination results support the suggestion that the ICP is normal. Volumetric analysis to establish ventricular volume norms (especially with aging), and regional analysis comparing intra- with extraventricular CSF compartments can help distinguish idiopathic NPH from atrophy. The definition of ventriculomegaly has not been standardized, and anatomical imaging is usually insufficient as a sole diagnostic test for hydrocephalus.

Injection of contrast material or radioactive tracer into CSF (cisternography) can allow detection of CSF flow distribution and clearance. Computed tomography scanning performed after injection of contrast into the CSF can identify sites of obstruction. Although radionuclide cisternography would appear to be an excellent method for detecting abnormal CSF circulation and clearance, its utility in the diagnosis of idiopathic NPH has been disappointing.^{17,20,125}

Magnetic resonance spectroscopy, functional MR imaging, positron emission tomography, and single-photon emission CT studies have been used for differential diagnosis and to assess the metabolic consequences of hydrocephalus. Although many reports implicate decreased CBF and even transient anaerobic metabolism,¹⁴³ these cannot be generalized to clinical practice.^{22,163,170,181} Based on recent research, some investigators suggest that MR imaging methods can determine ICP or measure CBF responses to treatment.⁷⁵ Clearly, advances in anatomical and functional imaging have the potential to allow more efficient initial diagnosis, treatment optimization, and recognition of failure.

Invasive Testing Methods. Contemporary invasive diagnostic procedures include CSF pressure monitoring (using lumbar puncture, spinal or ventricular catheter, and intracranial sensor), CSF infusion for measurement of R_{out} , lumbar CSF drainage, and shunt tapping for diagnosis of shunt failure. Because invasive techniques carry risk, they are often reserved for complex diagnostic problems, as in the identification of idiopathic NPH, chronic shunt failure, and shunt overdrainage, where the benefit of the invasive testing offsets the associated risks and burdens.

Monitoring of CSF pressure was first advocated as a diagnostic test for idiopathic NPH more than 30 years ago.^{138,141} Idiopathic NPH has the same abnormal CSF pressure waveforms originally described in brain tumor or acute injury.^{21,27,138} The Lundberg A-, B-, and C-waves are slow oscillations that reveal changes in the beat-to-beat amplitude and pulsatility of CSF pressure that are thought to reflect changes in cerebral blood volume associated with normal or abnormal cerebrovascular regulation. Eide et al.^{62,63,164} recently suggested that the amplitude of individual CSF pressure pulse waves may predict the response to CSF shunting in idiopathic NPH and in some children with hydrocephalus. Transfer function determination^{114,148} and analysis of phase shifts¹² have led to the proposal of a link between the CSF pressure pulsation and the overall system response, linking clinical monitoring with biomechanical modeling. Ultimately, invasive measures of CSF dynamics such as pressure waves, pulsatility, CSF space compliance, and absorption resistance may be critical to the identification and optimal treatment of hydrocephalus.

Biomarkers. Researchers using animal models have used CSF markers extensively,^{42,43,126} and so have investigators undertaking studies in humans.^{10,162} Although trends may be apparent in relative levels of catecholamines, energy metabolites, myelin basic protein, and glial fibrillary acidic protein, to name a few, the overall results of these studies are inconclusive because of variations in models and the complex clinical origins of the disorder. Tau protein and amyloid- β are being evaluated as biomarkers in idiopathic NPH.^{42,163} Although blood and CSF biomarkers based on CNS injury or metabolic or blood-brain barrier changes are unlikely to be specific to the diagnosis of hydrocephalus, they may still play an important role in determining the need to treat, the response to treatment, and also the prognosis.

Because of the difficulty in timely and sensitive identification of shunt infection, CSF markers of infection such as C-reactive protein,¹⁶⁰ if further developed, could play a large role in improving the treatment of shunt infection.

Neuropsychological Testing. Neuropsychological testing has long implicated problems in personality, cognition, and memory in children with hydrocephalus, and a subcortical dementia with short-term memory loss in adults with idiopathic NPH.^{26,56,149,172} Although neuropsychological evaluation is not specific for hydrocephalus, it can be used to assess response to treatment and determine prognosis. Treating idiopathic NPH results in substantially better neuropsychological improvement than previously thought, surpassing the modest improvement seen with pharmacological treatment of Alzheimer disease.¹⁷² Whether such cognitive improvement follows treatment of hydrocephalus in younger adults awaits demonstration.

Education. Because the majority of patients with hydro-

Priorities for hydrocephalus research

cephalus are children, most of the expertise in the diagnosis and treatment of this disorder has been among pediatric specialists. Nevertheless, there has been rapid growth in the number of adults with hydrocephalus for several reasons: children with the disorder survive to become adults; adults may experience secondary hydrocephalus from many causes; and idiopathic NPH is being recognized more frequently. Thus, if hydrocephalus continues to be perceived as “only” a pediatric disorder, it can be overlooked as a diagnostic consideration for adults.^{31,185} Therefore, education about hydrocephalus should be targeted to internists, geriatricians, family physicians, and neurologists and neurosurgeons who treat adults. Patient and family-oriented education and public awareness campaigns may help bring patients to the attention of physicians.

Current Treatment, Outcomes, and Complications

The causes, presentation, course of illness, and treatment paradigms for hydrocephalus are different for children and adults. Current treatment in both groups is based on a limited number of randomized controlled trials, case-control studies, prospective observational studies, and retrospective case series. Guidelines for the most common adult form of hydrocephalus (idiopathic NPH) have been developed,¹¹⁹ but paradoxically there are no guidelines for pediatric hydrocephalus, which is much more prevalent. Therefore, research needs related to treatment and outcomes of children may be broader. For example, treatment indications are not clearly defined, and there is significant variation between centers and surgeons in the decision to treat or observe. Thus, there is a need for objective methods to diagnose hydrocephalus and to select patients who will benefit from treatment.

There are no established criteria for selecting the most appropriate treatment (for example, shunt compared with ETV, type of valve, and technique of shunt insertion) for children of any age, although a randomized trial comparing shunt placement and ETV in infants (defined as children < 1 year old) is under way in Europe. Shunt obstruction and malfunction occur in approximately one third of children in the 1st year after shunt surgery.⁵⁴ Two randomized trials failed to show a difference in shunt failure rates among valve types.^{54,55,103}

There are few standardized outcome measures for either pediatric or adult hydrocephalus. Although shunt survival time and infection rates are commonly used, other important outcomes should be evaluated, such as neuropsychological and cognitive outcomes, which most likely are age-dependent. A Hydrocephalus Outcome Questionnaire has been developed and validated for children between 5 and 18 years of age.^{111,112}

Infection complicates 8 to 10% of shunt operations in children^{54,104,105,182} and may necessitate removal or externalization of the device, inpatient antibiotic therapy, and insertion of a new shunt. However, there are no standards for the diagnosis and treatment of shunt infection, and there is substantial variation in the duration of antibiotic therapy, decisions to remove or replace infected shunts, and length of hospital stay. The use of prophylactic antibiotics is generally accepted. Other strategies may reduce the risk of infection but require further study, such as antibiotic-impregnated shunt catheters,¹⁶¹ double gloving,^{112,147} and injection of

antibiotics into the shunt lumen during surgery.¹⁵⁰ Reinfection is alarmingly common (25%) and does not appear to be related to the duration of antibiotic therapy.¹⁰⁵ Shunts impregnated with rifampicin and clindamycin reduce reinfection from *Staphylococcus epidermidis*.⁹ Further studies of the effectiveness of antibiotic-impregnated external ventricular catheters and shunt catheters in the prevention and management of infection are warranted.

As discussed earlier, establishment of a hydrocephalus research infrastructure, including clinical data repositories, would facilitate longitudinal studies. The database should be broadly available to researchers and the physician community and should protect the privacy of patients, consistent with federal, local, and state regulations and laws.

Priorities for Future Treatments

The most forward-looking research priorities are for future treatments (Table 1). These include not only the development of novel medical devices, biomaterials, or surgical techniques, which represent a continuation of the first 50 years of hydrocephalus treatment research, but also the development of novel adjunctive (that is, pharmacological) therapies that should emerge from improved understanding of the basic biology of hydrocephalus and its impact on the brain.

Novel Medical Devices and Biomaterials

Shunts have been the fundamental treatment for hydrocephalus for nearly 50 years. Ironically, this may have caused many to believe that hydrocephalus was a “solved problem” that no longer required focused investigation.¹¹⁴ Yet, as described earlier, shunts are often associated with side effects and complications. Despite advances in their design, shunt failure nonetheless remains a significant problem. With the exception of ETV, which is used in a minority of patients, there are no alternative treatments and no proven medical or pharmacological treatments.

Medical devices and new treatments for hydrocephalus should be designed in accordance with contemporary understanding of the pathophysiological features of hydrocephalus. The goal is to bring new technologies and research discoveries together to improve the diagnosis and treatment of hydrocephalus for patients of all ages. This should include diagnostic aids that allow treatment to be individualized and optimized. For example, the decision to change the setting of an adjustable shunt is usually guided by clinical findings, subjective impressions, or neuroimaging, resulting in variations of practice. An objective optimization parameter is needed to guide such decisions. Possibilities include the analysis of CSF pulsation amplitude or MR imaging and transcranial Doppler methods.^{62,63,164}

There is no method to quantitate the pressure or flow function of shunts in real time in the outpatient setting. The development of devices that record these variables over time has the potential to improve short- and long-term treatment outcomes. Device size and inability to record ambulatory data have limited past efforts at telemetric ICP and shunt flow monitoring.^{39,70,187} Microelectromechanical versions of such devices, including the development of a smart monitor to allow query of shunt performance data, could overcome these limitations.

Devices that respond to changes in the internal milieu (for example, biochemical or physiological changes) may play an important role in future therapies. For example, it is likely that the functional characteristics of a shunt that are optimal for one set of conditions (posture, activity, heart rate, blood pressure, and so on) are suboptimal in another. Antisiphoning devices, which have been used to address postural effects for many years, are rudimentary, and improved physiological understanding is the key to developing improved devices. Last, although antibiotic-impregnated shunt components already exist, further research on biomaterials is needed to create shunt catheters with less risk of cellular occlusion or more resistance to bacterial adhesion, for example.

Adjunctive Therapies

Pharmacological agents (for example, acetazolamide) have been used as temporizing strategies to treat hydrocephalus in certain situations, especially in newborns, but drugs have never been considered a mainstay of treatment, nor have they been demonstrated to be effective in adults. Improved understanding of the basic science of injury and recovery mechanisms and homeostatic mechanisms such as control of cell volume, response of endothelial cells to pulsatile shear, CSF absorption or secretion processes, and genetic regulation has great potential to lead to the development of pharmacological or biological agents to prevent or treat hydrocephalus and enhance clinical recovery. There is a need to determine the potential role of stem cell therapy, for example, to supply trophic agents or promote myelin formation. One model worth considering is the Stroke Therapy Academic Industry Roundtable, which presents recommendations for preclinical neuroprotective and restorative drug development.²

Human Research: Ethical Considerations

Considering that translational research in hydrocephalus represents exciting and unexplored territory, it will be vital that collaboration between basic and clinical scientists occurs to assure that promising laboratory methods transfer to humans. In addition, recognizing that hydrocephalus most often affects persons considered vulnerable human research candidates (infants, children, and adults with dementia and impaired decision-making capacity), and that neither the immediate nor long-term effects of novel pharmacological or biological agents can be predicted, it will be important to consider the ethical boundaries and safeguards of such research prospectively.

Acknowledgments

We thank Susan Moichan, Ph.D., for her contributions to this study. We greatly appreciate the collaboration with the professionals at the NIH who helped organize, promote, and fund this workshop. We are most indebted to the efforts of the leadership of the Hydrocephalus Association, especially Emily Fudge, Dory Kranz, and Pip Marks. These individuals were instrumental in initiating discussions with the NIH that led to the workshop. We also thank Curt Stewart of the BrainChild Foundation for his generous financial support of an organizational meeting in Baltimore in October 2004 that led to the workshop proposal to the National Institute of Neurological Disorders and Stroke. Yandong Zhou was instrumental in formulating the bibliography.

Appendix 1

Breakout sessions and key questions

Breakout Session I: Injury Mechanisms, Neuroprotection, and Cellular Recovery in Hydrocephalus

Breakout Session II: New Insights into the Pathophysiology of CSF Circulation—Implications of Shunt Design or Other Treatments for Hydrocephalus

Breakout Session III: Clinical Tools, Clinical Trials, and Clinical Outcomes

Key questions for all breakout sessions:

1. What is the critical information that has been missing or needs to be reconsidered to advance our understanding of the pathophysiological features of hydrocephalus, its clinical manifestations, its treatment, and its outcomes?
2. What is the best way to obtain this information?
3. How can research in related disorders be used to improve our understanding of hydrocephalus, and vice versa?
4. How can clinicians, clinical researchers, and basic scientists collaborate more effectively?
5. What are the critical differences between pediatric and adult hydrocephalus?

Appendix 2

Participants and contributors

Workshop Leadership Committee:

Chair, James P. “Pat” McAllister, Wayne State University, Detroit, Michigan; *Chair*, Marion L. “Jack” Walker, University of Utah, Salt Lake City, Utah; *Chair*, Michael A. Williams, Johns Hopkins Medical Institutions, Baltimore, Maryland; Dory Kranz, Hydrocephalus Association, San Francisco, California; Katrina Gwinn, NINDS, Bethesda, Maryland

Breakout Session 1—Injury Mechanisms, Neuroprotection, and Cellular Recovery in Hydrocephalus

Facilitators: James P. “Pat” McAllister, Wayne State University, Detroit, Michigan; Michael A. Williams, Johns Hopkins Medical Institutions, Baltimore, Maryland; Hazel Jones, University of Florida, Gainesville, Florida

Breakout Session 2—New Insights into the Pathophysiology of CSF Circulation: Implications of Shunt Design or Other Treatments for Hydrocephalus

Facilitators: Marion L. “Jack” Walker, University of Utah, Salt Lake City, Utah; Michael Egnor, State University of New York, Stony Brook, New York; Marvin L. Sussman, Miami, Florida

Breakout Session 3—Clinical Tools, Clinical Trials, and Clinical Outcomes

Facilitators: Anthony Marmarou, Virginia Commonwealth University, Richmond, Virginia; John Kestle, University of Utah, Salt Lake City, Utah; Dory Kranz, Hydrocephalus Association, San Francisco, California

Additional invited moderators and speakers:

Marvin Bergsneider, University of California Los Angeles, California; Peter Black, Brigham and Women’s Hospital, Boston, Massachusetts; Marc Del Bigio, University of Manitoba, Winnipeg, Canada; Maureen Dennis, University of Toronto, Canada; James Drake, University of Toronto, Canada; David Frim, University of Chicago, Illinois; Neill Graff-Radford, Mayo Clinic, Jacksonville, Florida; Stephen Graft, NIH Office of Rare Diseases, Bethesda, Maryland; Bermans Iskandar, University of Wisconsin, Madison, Wisconsin; Simon John, Howard Hughes Medical Institute, Bar Harbor, Maine; Miles Johnston, Sunnybrook & Women’s College, Toronto, Canada; Dianne Kean, Seeking Techniques Advancing Research in Shunts, Detroit, Michigan; Mark Luciano, Cleveland Clinic Foundation, Cleveland, Ohio; Joseph Madsen, Harvard Medical School, Boston, Massachusetts; Ed Oidfield, NINDS, Bethesda, Maryland; Harold Rekaté, Barrow Neurological Institute, Phoenix, Arizona; M. Elizabeth Ross, Weill Medical College of Cornell Uni-

Priorities for hydrocephalus research

versity, New York, New York; Gerald Silverberg, Stanford University, Stanford, California; Marcy Speer, Duke University, Durham, North Carolina; Henning Stolze, Universitätsklinikum Schleswig-Holstein Campus, Kiel, Germany

Invited participants:

Duane Alexander, National Institute of Child Health and Human Development, Bethesda, Maryland; Anthony Avellino, University of Washington, Seattle, Washington; Roger Bayston, University of Nottingham, United Kingdom; Timothy Brei, Indiana University, Indianapolis, Indiana; Cindy Brownstein, Spina Bifida Association of America, Washington, DC; Neil Buckholtz, National Institute on Aging, Bethesda, Maryland; Debby Buffa, Hydrocephalus Support Group, St. Louis, Missouri; Stephanie Clipper, NIH, Bethesda, Maryland; Adolfo Correa, Centers for Disease Control, Atlanta, Georgia; Denise Dougherty, Agency for Healthcare Research and Quality, Rockville, Maryland; Marion Emr, NIH, Bethesda, Maryland; Heather Etchevers, Hôpital Necker-Enfants Malades, Paris, France; Rebecca Farkas, NINDS, Bethesda, Maryland; John Ferguson, NIH, Bethesda, Maryland; Debbi Fields, National Hydrocephalus Foundation, Lakewood, California; Helen Fillmore, Virginia Commonwealth University, Richmond, Virginia; Emily Fudge, Hydrocephalus Association, San Francisco, California; Dan Greitz, Karolinska University Hospital, Stockholm, Sweden; John Heiss, NINDS, Bethesda, Maryland; Argye Hillis, Johns Hopkins Medical Institutions, Baltimore, Maryland; Maya House, Spina Bifida Association of America, Washington, DC; Debra Jaurigue, University of Vermont, Rutland; Conrad Johanson, Brown University, Providence, Rhode Island; Petra Klinge, International Neuroscience Institute, Hannover, Germany; Story Landis, NINDS, Bethesda, Maryland; Karen Langston, Oviedo, Florida; Vance Lemmon, University of Miami, Florida; Ellen Liberman, National Eye Institute, Bethesda, Maryland; Robert McAllister, Intracranial Hypertension Research Foundation, Vancouver, Washington; Dave McLone, Children's Memorial Hospital, Chicago, Illinois; Susan Molchan, National Institute on Aging, Bethesda, Maryland; Marcelle Morrison-Bogorad, National Institute on Aging, Bethesda, Maryland; Mary Lou Oster-Granite, National Institute of Child Health and Human Development, Bethesda, Maryland; Joe Panerazio, NINDS, Bethesda, Maryland; Carlos Pena, Food and Drug Administration, Rockville, Maryland; Grace Peng, National Institute of Biomedical Imaging and BioEngineering, Bethesda, Maryland; Anna Penn, Stanford University, Stanford, California; Richard Penn, University of Chicago, Illinois; Victor Perry, University of California San Francisco, California; Dorothy Poppe, American Syringomyelia Alliance Project, Staten Island, New York; Paul Rosenberg, Johns Hopkins Medical Institutions, Baltimore, Maryland; Hilary Rossen, Northeast Ohio Hydrocephalus Support Group, Cleveland, Ohio; Paul Scott, NINDS, Bethesda, Maryland; Siv Sivaloganathan, University of Waterloo, Canada; Mary Smellie-Decker, Children's Hospital of Michigan, Detroit, Michigan; Giovanna Spinella, NIH, Bethesda, Maryland; Curt Stewart, BrainChild Foundation, Carefree, Arizona; Greg Tocco, Hydrocephalus Foundation, Inc., Saugus, Massachusetts; Mark Wagshul, State University of New York, Stony Brook, New York; Bob Zalutsky, NINDS, Bethesda, Maryland

References

1. Albeck MJ, Børgesen SE, Gjerris F, Schmidt JF, Sorensen PS: Intracranial pressure and cerebrospinal fluid outflow conductance in healthy subjects. **J Neurosurg** 74:597–600, 1991
2. Anonymous: Recommendations for standards regarding preclinical neuroprotective and restorative drug development. **Stroke** 30:2752–2758, 1999
3. Badano JL, Mitsuma N, Beales PL, Katsanis N: The ciliopathies: an emerging class of human genetic disorders. **Annu Rev Genomics Hum Genet** 7:125–148, 2006
4. Balasubramaniam J, Del Bigio MR: Analysis of age-dependant alteration in the brain gene expression profile following induction of hydrocephalus in rats. **Exp Neurol** 173:105–113, 2002
5. Baru JS, Bloom DA, Muraszko K, Koop CE: John Holter's shunt. **J Am Coll Surg** 192:79–85, 2001
6. Bateman GA: The reversibility of reduced cortical vein compli-

ance in normal-pressure hydrocephalus following shunt insertion. **Neuroradiology** 45:65–70, 2003

7. Bateman GA, Levi CR, Schofield P, Wang Y, Lovett EC: The pathophysiology of the aqueduct stroke volume in normal pressure hydrocephalus: can co-morbidity with other forms of dementia be excluded? **Neuroradiology** 47:741–748, 2005
8. Bátiz LF, Páez P, Jiménez AJ, Rodríguez S, Wagner C, Pérez-Figares JM, et al: Heterogeneous expression of hydrocephalic phenotype in the hyh mice carrying a point mutation in alpha-SNAP. **Neurobiol Disease** 23:152–168, 2006
9. Bayston R, Ashraf W, Bhundia C: Mode of action of an antimicrobial biomaterial for use in hydrocephalus shunts. **J Antimicrob Chemother** 53:778–782, 2004
10. Beems T, Simons KS, Van Geel WJ, De Reus HP, Vos PE, Verbeek MM: Serum- and CSF-concentrations of brain specific proteins in hydrocephalus. **Acta Neurochir (Wien)** 145:37–43, 2003
11. Bergsneider M: Evolving concepts of cerebrospinal fluid physiology. **Neurosurg Clin N Am** 12:631–638, 2001
12. Bergsneider M: Hydrocephalus: new theories and new shunts? **Clin Neurosurg** 52:120–126, 2005
13. Bergsneider M, Alwan AA, Falkson L, Rubinstein EH: The relationship of pulsatile cerebrospinal fluid flow to cerebral blood flow and intracranial pressure: a new theoretical model. **Acta Neurochir Suppl** 71:266–268, 1998
14. Bering EA: Circulation of the cerebrospinal fluid. Demonstration of the choroid plexuses as the generator of the force for flow of fluid and ventricular enlargement. **J Neurosurg** 19:405–413, 1962
15. Bering EA Jr: Choroid plexus and arterial pulsation of cerebrospinal fluid: demonstration of the choroid plexuses as a cerebrospinal fluid pump. **AMA Arch Neurol Psychiatry** 73:165–172, 1955
16. Bering EA, Ingraham FD: The arterial pulsation of the cerebrospinal fluid; its origin, configuration and possible clinical importance. **Trans Am Neurol Assoc** 3:44–52, 1953
17. Black PM: Idiopathic normal-pressure hydrocephalus. Results of shunting in 62 patients. **J Neurosurg** 52:371–377, 1980
18. Bloch O, Auguste KI, Manley GT, Verkman AS: Accelerated progression of kaolin-induced hydrocephalus in aquaporin-4-deficient mice. **J Cereb Blood Flow Metab** 26:1527–1537, 2006
19. Boillat CA, Jones HC, Kaiser GL, Harris NG: Ultrastructural changes in the deep cortical pyramidal cells of infant rats with inherited hydrocephalus and the effect of shunt treatment. **Exp Neurol** 147:377–388, 1997
20. Børgesen SE: Conductance to outflow of CSF in normal pressure hydrocephalus. **Acta Neurochir (Wien)** 71:1–45, 1984
21. Børgesen SE, Gjerris F: The predictive value of conductance to outflow of CSF in normal pressure hydrocephalus. **Brain** 105: 65–86, 1982
22. Bradley WG: Cerebrospinal fluid dynamics and shunt responsiveness in patients with normal-pressure hydrocephalus. **Mayo Clin Proc** 77:507–508, 2002
23. Braun KPI, Vandertop WP, Gooskens RH, Tulleken KA, Nicolay K: NMR spectroscopic evaluation of cerebral metabolism in hydrocephalus: a review. **Neurol Res** 22:51–64, 2000
24. Bruni JE, Del Bigio MR, Cardoso ER, Persaud TV: Hereditary hydrocephalus in laboratory animals and humans. **Exp Pathol** 35:239–246, 1988
25. Carter BJ, Morel L, Jones HC: Characterization of inherited hydrocephalus in the LEW/Jms rat. **Society for Neuroscience Program**: No. 311.317, 2002 (Abstract)
26. Chang S, Agarwal S, Williams MA, Rigamonti D, Hillis AE: Demographic factors influence cognitive recovery after shunt for normal-pressure hydrocephalus. **Neurologist** 12:39–42, 2006
27. Chawla JC, Hulme A, Cooper R: Intracranial pressure in patients with dementia and communicating hydrocephalus. **J Neurosurg** 40:376–380, 1974
28. Chopp M, Portnoy HD: Systems analysis of intracranial pressure. Comparison with volume-pressure test and CSF-pulse amplitude analysis. **J Neurosurg** 53:516–527, 1980

29. Chopp M, Portnoy HD, Branch C: Hydraulic model of the cerebrovascular bed: an aid to understanding the volume-pressure test. **Neurosurgery** 13:5–11, 1983
30. Chovanes GI, McAllister JP, Lamperti AA, Salotto AG, Truex RC: Monoamine alterations during experimental hydrocephalus in neonatal rats. **Neurosurgery** 22:86–91, 1988
31. Cowan JA, McGirt MJ, Woodworth G, Rigamonti D, Williams MA: The syndrome of hydrocephalus in young and middle-aged adults (SHYMA). **Neurol Res** 27:540–547, 2005
32. Crews L, Wyss-Coray T, Masliah E: Insights into the pathogenesis of hydrocephalus from transgenic and experimental animal models. **Brain Pathol** 14:312–316, 2004
33. Czosnyka M: Pulsatility index. **J Neurosurg** 94:685–686, 2001
34. Czosnyka M, Czosnyka ZH, Whitfield PC, Donovan T, Pickard JD: Age dependence of cerebrospinal pressure-volume compensation in patients with hydrocephalus. **J Neurosurg** 94:482–486, 2001
35. Czosnyka M, Maksymowicz W, Batorski L, Koszewski W, Czosnyka Z: Comparison between classic-differential and automatic shunt functioning on the basis of infusion tests. **Acta Neurochir (Wien)** 106:1–8, 1990
36. Czosnyka M, Piechnik S, Koszewski W, Laniewski P, Maksymowicz W, Paluszek K, et al: The dynamics of cerebral blood flow perfusion pressure and CSF circulation—a modelling study, in Avezaat CJJ (ed): **Intracranial Pressure VIII**. Berlin: Springer-Verlag, 1993, pp 699–706
37. Czosnyka M, Whitehouse H, Smielewski P, Simac S, Pickard JD: Testing of cerebrospinal compensatory reserve in shunted and non-shunted patients: a guide to interpretation based on an observational study. **J Neurol Neurosurg Psychiatry** 60:549–558, 1996
38. Dandy WE: Ventriculography following injection of air into the cerebral ventricles. **Ann Surg** 68:4–11, 1918
39. de Jong DA, Maas AI, den Ouden AH, de Lange SA: Long-term intracranial pressure monitoring. **Med Prog Technol** 10:89–96, 1983
40. Del Bigio MR: Calcium-mediated proteolytic damage in white matter of hydrocephalic rats? **J Neuropathol Exp Neurol** 59:946–954, 2000
41. Del Bigio MR: Cellular damage and prevention in childhood hydrocephalus. **Brain Pathol** 14:317–324, 2004
42. Del Bigio MR: Hydrocephalus-induced changes in the composition of cerebrospinal fluid. **Neurosurgery** 25:416–423, 1989
43. Del Bigio MR: Neuropathological changes caused by hydrocephalus. **Acta Neuropathol** 85:573–585, 1993
44. Del Bigio MR: Pathophysiologic consequences of hydrocephalus. **Neurosurg Clin N Am** 12:639–649, 2001
45. Del Bigio MR, Cardoso ER, Halliday WC: Neuropathological changes in chronic adult hydrocephalus: cortical biopsies and autopsy findings. **Can J Neurol Sci** 24:121–126, 1997
46. Del Bigio MR, da Silva MC, Drake JM, Tuor UI: Acute and chronic cerebral white matter damage in neonatal hydrocephalus. **Can J Neurol Sci** 21:299–305, 1994
47. Del Bigio MR, Kanfer JN, Zhang YW: Myelination delay in the cerebral white matter of immature rats with kaolin-induced hydrocephalus is reversible. **J Neuropathol Exp Neurol** 56:1053–1066, 1997
48. Del Bigio MR, McAllister JP II: Hydrocephalus—pathology, in Choux M, DiRocco R, Hockley AD, et al (eds): **Pediatric Neurosurgery**. Philadelphia: Saunders, 1999, pp 217–236
49. Del Bigio MR, Vriend JP: Monoamine neurotransmitters and amino acids in the cerebrum and striatum of immature rats with kaolin-induced hydrocephalus. **Brain Res** 798:119–126, 1998
50. Del Bigio MR, Wilson MJ, Enno T: Chronic hydrocephalus in rats and humans: white matter loss and behavior changes. **Ann Neurol** 53:337–346, 2003
51. Di Rocco C, Di Trapani G, Pettorossi VE, Caldarelli M: On the pathology of experimental hydrocephalus induced by artificial increase in endoventricular CSF pulse pressure. **Childs Brain** 5:81–95, 1979
52. Di Rocco C, Pettorossi VE, Caldarelli M, Mancinelli R, Velardi F: Communicating hydrocephalus induced by mechanically increased amplitude of the intraventricular cerebrospinal fluid pressure: experimental studies. **Exp Neurol** 59:40–52, 1978
53. Dombrowski SM, Schenk S, Leichter A, Leibson Z, Fukumachi K, Luciano MG: Chronic hydrocephalus-induced changes in cerebral blood flow: mediation through cardiac effects. **J Cereb Blood Flow Metab** 26:1298–1310, 2006
54. Drake JM, Kestle JR, Milner R, Cinalli G, Boop F, Piatt J, et al: Randomized trial of cerebrospinal fluid shunt valve design in pediatric hydrocephalus. **Neurosurgery** 43:294–303, 1998
55. Drake JM, Kestle JT: Determining the best cerebrospinal fluid shunt valve design: the pediatric valve design trial. **Neurosurgery** 43:1259–1260, 1998
56. Duinkerke A, Williams MA, Rigamonti D, Hillis AE: Cognitive recovery in idiopathic normal pressure hydrocephalus after shunt. **Cogn Behav Neurol** 17:179–184, 2004
57. Edwards MS, Harrison MR, Halks-Miller M, Nakayama DK, Berger MS, Glick PL, et al: Kaolin-induced congenital hydrocephalus in utero in fetal lambs and rhesus monkeys. **J Neurosurg** 60:115–122, 1984
58. Edwards RJ, Dombrowski SM, Luciano MG, Pople IK: Chronic hydrocephalus in adults. **Brain Pathol** 14:325–336, 2004
59. Egnor M, Rosiello A, Zheng L: A model of intracranial pulsations. **Pediatr Neurosurg** 35:284–298, 2001
60. Egnor M, Wagshul M, Zheng L, Rosiello A: Resonance and the synchrony of arterial and CSF pulsations. **Pediatr Neurosurg** 38:273–276, 2003
61. Egnor M, Zheng L, Rosiello A, Gutman F, Davis R: A model of pulsations in communicating hydrocephalus. **Pediatr Neurosurg** 36:281–303, 2002
62. Eide PK: Assessment of childhood intracranial pressure recordings using a new method of processing intracranial pressure signals. **Pediatr Neurosurg** 41:122–130, 2005
63. Eide PK: Intracranial pressure parameters in idiopathic normal pressure hydrocephalus patients treated with ventriculo-peritoneal shunts. **Acta Neurochir (Wien)** 148:21–29, 2006
64. Eide PK, Due-Tonnessen B, Helseth E, Lundar T: Differences in quantitative characteristics of intracranial pressure in hydrocephalic children treated surgically or conservatively. **Pediatr Neurosurg** 36:304–313, 2002
65. Eide PK, Helseth E, Due-Tonnessen B, Lundar T: Changes in intracranial pressure after calvarial expansion surgery in children with slit ventricle syndrome. **Pediatr Neurosurg** 35:195–204, 2001
66. Ekstedt J: CSF hydrodynamic studies in man. 1. Method of constant pressure CSF infusion. **J Neurol Neurosurg Psychiatry** 40:105–119, 1977
67. Ekstedt J: CSF hydrodynamic studies in man. 2. Constant pressure CSF infusion. **J Neurol Neurosurg Psychiatry** 41:345–353, 1977
68. Eskandari R, McAllister JP II, Miller JM, Ding Y, Ham SD, Shearer DM, et al: Effects of hydrocephalus and ventriculoperitoneal shunt therapy on afferent and efferent connections in the feline sensorimotor cortex. **J Neurosurg** 101 (2 Suppl):196–210, 2004
69. Fransen E, D'Hooge R, Van Camp G, Verhoye M, Sijbers J, Reyniers E, et al: L1 knockout mice show dilated ventricles, vermiform hypoplasia and impaired exploration patterns. **Hum Mol Genet** 7:999–1009, 1998
70. Frim DM, Lathrop D: Telemetric assessment of intracranial pressure changes consequent to manipulations of the Codman-Medos programmable shunt valve. **Pediatr Neurosurg** 33:237–242, 2000
71. Fukuhara T, Luciano MG: Clinical features of late-onset idiopathic aqueductal stenosis. **Surg Neurol** 55:132–137, 2001
72. Galbreath E, Kim SJ, Park K, Brenner M, Messing A: Overexpression of TGF-beta 1 in the central nervous system of transgenic mice results in hydrocephalus. **J Neuropathol Exp Neurol** 54:339–349, 1995

Priorities for hydrocephalus research

73. Garton HJ, Piatt JH Jr: Hydrocephalus. **Pediatr Clin North Am** **51**:305–325, 2004
74. Glick PL, Harrison MR, Halks-Miller M, Adzick NS, Nakayama DK, Anderson JH, et al: Correction of congenital hydrocephalus in utero II: efficacy of in utero shunting. **J Pediatr Surg** **19**: 870–881, 1984
75. González RG, Fischman AJ, Guimaraes AR, Carr CA, Stern CE, Halpern EF, et al: Functional MR in the evaluation of dementia: correlation of abnormal dynamic cerebral blood volume measurements with changes in cerebral metabolism on positron emission tomography with fludeoxyglucose F 18. **AJNR Am J Neuroradiol** **16**:1763–1770, 1995
76. Gooskens RH, Willemse J, Gielen C: Cerebrospinal fluid dynamics and cerebrospinal fluid infusion in children. Part I: a revised method and a review. **Neuropediatrics** **16**:115–120, 1985
77. Greitz D: Radiological assessment of hydrocephalus: new theories and implications for therapy. **Neurosurg Rev** **27**:145–167, 2004
78. Greitz D, Franck A, Nordell B: On the pulsatile nature of intracranial and spinal CSF-circulation demonstrated by MR imaging. **Acta Radiol** **34**:321–328, 1993
79. Greitz D, Wirestam R, Franck A, Nordell B, Thomsen C, Stahlberg F: Pulsatile brain movement and associated hydrodynamics studied by magnetic resonance phase imaging. The Monro-Kellie doctrine revisited. **Neuroradiology** **34**:370–380, 1992
80. Gupta N, Park J, Solomon C, Kranz D, Wrensch M, Wu Y: Long-term outcomes in patients with treated childhood hydrocephalus. **J Neurosurg** **106** (5 Suppl):334–339, 2007
81. Hakim S: **Some Observations on C.S.F. Pressure. Hydrocephalic Syndrome in Adults with “Normal” C.S.F. Pressure. (Recognition of a New Syndrome). Thesis No. 957.** Bogota, Columbia: Javeriana University School of Medicine, 1964
82. Hakim S, Adams RD: The special clinical problem of symptomatic hydrocephalus with normal cerebrospinal fluid pressure. Observations on cerebrospinal fluid hydrodynamics. **J Neurol Sci** **2**:307–327, 1965
83. Hakim S, Venegas JG, Burton JD: The physics of the cranial cavity, hydrocephalus and normal pressure hydrocephalus: mechanical interpretation and mathematical model. **Surg Neurol** **5**:187–210, 1976
84. Hanigan WC, Morgan AM, Anderson RJ, Bradle P, Cohen HS, Cusack TJ, et al: Incidence and neurodevelopmental outcome of periventricular hemorrhage and hydrocephalus in a regional population of very low birth weight infants. **Neurosurgery** **29**: 701–706, 1991
85. Harris NG, Jones HC, Patel S: Ventricle shunting in young H-Tx rats with inherited congenital hydrocephalus: a quantitative histological study of cortical grey matter. **Childs Nerv Syst** **10**: 293–301, 1994
86. Harris NG, McAllister JP II, Conaughty JM, Jones HC: The effect of inherited hydrocephalus and shunt treatment on cortical pyramidal cell dendrites in the infant H-Tx rat. **Exp Neurol** **141**:269–279, 1996
87. Harris NG, Plant HD, Inglis BA, Briggs RW, Jones HC: Neurochemical changes in the cerebral cortex of treated and untreated hydrocephalic rat pups quantified with in vitro ¹H-NMR spectroscopy. **J Neurochem** **68**:305–312, 1997
88. Hochwald GM: Animal models of hydrocephalus: recent developments. **Proc Soc Exp Biol Med** **178**:1–11, 1985
89. Hoffman PL, Walter R, Bulat M: An enzymatically stable peptide with activity in the central nervous system: its penetration through the blood-CSF barrier. **Brain Res** **122**:87–94, 1977
90. Hu X, Alwan AA, Rubinstein EH, Bergsneider M: Reduction of compartment compliance increases venous flow pulsatility and lowers apparent vascular compliance: implications for cerebral blood flow hemodynamics. **Med Eng Phys** **28**:304–314, 2006
91. Ishizaki R, Tashiro Y, Inomoto T, Hashimoto N: Acute and subacute hydrocephalus in a rat neonatal model: correlation with functional injury of neurotransmitter systems. **Pediatr Neurosurg** **33**: 298–305, 2000
92. Johanson C, Del Bigio MR, Kinsman S, Miyani J, Pattisapu JV, Robinson M, et al: New models for analysing hydrocephalus and disorders of CSF volume transmission. **Br J Neurosurg** **15**: 281–283, 2001
93. Johanson C, McMillan P, Tavares R, Spangenberg A, Duncan J, Silverberg G, et al: Homeostatic capabilities of the choroid plexus epithelium in Alzheimer’s disease. **Cerebrospinal Fluid Res** **1**:3, 2004
94. Johanson CE, Flaherty S, Duncan J, Stopa EG, Silverberg GD: Aging rat brain: a model for analyzing interactions among CSF dynamics, ventriculomegaly, and the β -amyloid retention in Alzheimer’s disease. **Cerebrospinal Fluid Res** **2** (1 Suppl):S6, 2005
95. Jones HC, Carter BJ, Depelteau JS, Roman M, Morel L: Chromosomal linkage associated with disease severity in the hydrocephalic H-Tx rat. **Behav Genet** **31**:101–111, 2001
96. Jones HC, Carter BJ, Morel L: Characteristics of hydrocephalus expression in the LEW/Jms rat strain with inherited disease. **Childs Nerv Syst** **19**:11–18, 2003
97. Jones HC, Chen GF, Yehia BR, Carter BJ, Akins EJ, Wolpin LC: Single and multiple congenic strains for hydrocephalus in the H-Tx rat. **Mamm Genome** **16**:251–261, 2005
98. Jones HC, Depelteau JS, Carter BJ, Somera KC: The frequency of inherited hydrocephalus is influenced by intrauterine factors in H-Tx rats. **Exp Neurol** **176**:213–220, 2002
99. Jones HC, Harris NG, Rocca JR, Andersohn RW: Progressive changes in cortical metabolites at three stages of infantile hydrocephalus studied by in vitro NMR spectroscopy. **J Neurotrauma** **14**:587–602, 1997
100. Jones HC, Totten CF, Mayorga DA, Yue M, Carter BJ: Genetic loci for ventricular dilatation in the LEW/Jms rat with fetal-onset hydrocephalus are influenced by gender and genetic background. **Cerebrospinal Fluid Res** **2**:2, 2005
101. Jones HC, Yehia B, Chen GF, Carter BJ: Genetic analysis of inherited hydrocephalus in a rat model. **Exp Neurol** **190**:79–90, 2004
102. Kellie G: An account of the appearances observed in the dissection of two of the three individuals presumed to have perished in the storm of the 3rd, and whose bodies were discovered in the vicinity of Leith on the morning of the 4th November 1821 with some reflections on the pathology of the brain. **The Transactions of the Medico-Chirurgical Society of Edinburgh** **1**:84–169, 1824
103. Kestle J, Drake J, Milner R, Sainte-Rose C, Cinalli G, Boop F, et al: Long-term follow-up data from the Shunt Design Trial. **Pediatr Neurosurg** **33**:230–236, 2000
104. Kestle JR: Pediatric hydrocephalus: current management. **Neurol Clin** **21**:883–895, 2003
105. Kestle JR, Garton HJ, Whitehead WE, Drake JM, Kulkarni AV, Cochrane DD, et al: Management of shunt infections: a multicenter pilot study. **J Neurosurg** **105** (3 Suppl):177–181, 2006
106. Khan OH, Enno TL, Del Bigio MR: Brain damage in neonatal rats following kaolin induction of hydrocephalus. **Exp Neurol** **200**:311–320, 2006
107. Kiefer M, Eymann R, Steudel WI: The dynamic infusion test in rats. **Childs Nerv Syst** **16**:451–456, 2000
108. Kosteljanetz M: Resistance to outflow of cerebrospinal fluid determined by bolus injection technique and constant rate steady state infusion in humans. **Neurosurgery** **16**:336–340, 1985
109. Kriebel RM, McAllister JP: Pathology of the hippocampus in experimental feline infantile hydrocephalus. **Neurol Res** **22**: 29–36, 2000
110. Kriebel RM, Shah AB, McAllister JP: The microstructure of cortical neuropil before and after decompression in experimental infantile hydrocephalus. **Exp Neurol** **119**:89–98, 1993
111. Kulkarni AV: Distribution-based and anchor-based approaches provided different interpretability estimates for the Hydrocephalus Outcome Questionnaire. **J Clin Epidemiol** **59**:176–184, 2006

112. Kulkarni AV, Drake JM, Lamberti-Pasculli M: Cerebrospinal fluid shunt infection: a prospective study of risk factors. **J Neurosurg** **94**:195–201, 2001
113. Lovely TJ, McAllister JP, Miller DW, Lamperti AA, Wolfson BJ: Effects of hydrocephalus and surgical decompression on cortical norepinephrine levels in neonatal cats. **Neurosurgery** **24**:43–52, 1989
114. Madsen JR, Egnor M, Zou R: Cerebrospinal fluid pulsatility and hydrocephalus: the fourth circulation. **Clin Neurosurg** **53**:48–52, 2006
115. Madsen JR, Zou R, Egnor MR, Luciano MG, Dombrowski S, McCormack EJ, et al: Variation in the notch filter response of the ICP to arterial pulse pressure. **AANS Abstract ID 29935**, 2005
116. Mangano FT, McAllister JP, Jones HC, Johnson MJ, Kriebel RM: The microglial response to progressive hydrocephalus in a model of inherited aqueductal stenosis. **Neurol Res** **20**:697–704, 1998
117. Mao X, Enno TL, Del Bigio MR: Aquaporin 4 changes in rat brain with severe hydrocephalus. **Eur J Neurosci** **23**:2929–2936, 2006
118. Marmarou A, Bergsneider M, Klinge P, Relkin N, Black PM: The value of supplemental prognostic tests for the preoperative assessment of idiopathic normal-pressure hydrocephalus. **Neurosurgery** **57** (3 Suppl):S17–S28, 2005
119. Marmarou A, Bergsneider M, Relkin N, Klinge P, Black PM: Development of guidelines for idiopathic normal-pressure hydrocephalus: introduction. **Neurosurgery** **57** (3 Suppl):S1–S3, 2005
120. Marmarou A, Black P, Bergsneider M, Klinge P, Relkin N, International NPH Consultant Group: Guidelines for management of idiopathic normal pressure hydrocephalus: progress to date. **Acta Neurochir Suppl** **95**:237–240, 2005
121. Marmarou A, Young HF, Aygok GA, Sawauchi S, Tsuji O, Yamamoto T, et al: Diagnosis and management of idiopathic normal-pressure hydrocephalus: a prospective study in 151 patients. **J Neurosurg** **102**:987–997, 2005
122. Matsumoto S, Hirayama A, Yamasaki S, Shirataki K, Fujiwara K: Comparative study of various models of experimental hydrocephalus. **Childs Brain** **1**:236–242, 1975
123. McAllister JP, Chovan P: Neonatal hydrocephalus. Mechanisms and consequences. **Neurosurg Clin N Am** **9**:73–93, 1998
124. McAllister JP, Maugans TA, Shah MV, Trux RC: Neuronal effects of experimentally induced hydrocephalus in newborn rats. **J Neurosurg** **63**:776–783, 1985
125. McCullough DC, Harbert JC, Di Chiro G, Ommaya AK: Prognostic criteria for cerebrospinal fluid shunting from isotope cisternography in communicating hydrocephalus. **Neurology** **20**:594–598, 1970
126. McGirt MJ, Woodworth G, Coon AL, Thomas G, Williams MA, Rigamonti D: Diagnosis, treatment, and analysis of long-term outcomes in idiopathic normal-pressure hydrocephalus. **Neurosurgery** **57**:699–705, 2005
127. Meier U, Bartels P: The importance of the intrathecal infusion test in the diagnosis of normal pressure hydrocephalus. **J Clin Neurosci** **9**:260–267, 2002
128. Meier U, Kiefer M, Bartels P: The ICP-dependency of resistance to cerebrospinal fluid outflow: a new mathematical method for CSF-parameter calculation in a model with H-TX rats. **J Clin Neurosci** **9**:58–63, 2002
129. Meier U, Künzel B, Zeilinger FS, Riederer A: [Pressure-dependent flow resistance in craniopsinal cerebrospinal fluid dynamics: a calculation model for diagnosis of normal pressure hydrocephalus.] **Biomed Tech (Berl)** **45**:26–33, 2000 (Ger)
130. Meuli M, Meuli-Simmen C, Yingling CD, Hutchins GM, Timmel GB, Harrison MR, et al: In utero repair of experimental myelomeningocele saves neurological function at birth. **J Pediatr Surg** **31**:397–402, 1996
131. Miller JM, Kumar R, McAllister JP, Krause GS: Gene expression analysis of the development of congenital hydrocephalus in the H-Tx rat. **Brain Res** **1075**:36–47, 2006
132. Miyake H, Eghwrudjakpor P, Sakamoto T, Kurisaka M, Mori K, Matsumoto S, et al: Neurotransmitter changes in hydrocephalus: effects of cerebral metabolic activator on kaolin-induced hydrocephalus, in Matsumoto S, Tamaki N (eds): **Hydrocephalus: Pathogenesis and Treatment**. Tokyo: Springer-Verlag, 1991, pp 68–74
133. Miyan J, Nabiyouni M, Zendah M: Development of the brain: a vital role for cerebrospinal fluid. **Can J Physiol Pharmacol** **81**:317–328, 2003
134. Miyan JA, Khan MI, Kawarada Y, Sugiyama T, Bannister CM: Cell death in the brain of the HTx rat. **Eur J Pediatr Surg** **8**:43–48, 1998
135. Monro A: **Observations on the Structure and Function of the Nervous System**. Edinburgh: Creech & Johnson, 1823
136. Morgan FW, Stewart JA, Smith AN, Tarnuzzer RW: Differential expression of stress response genes in the H-Tx rat model of congenital hydrocephalus. **Brain Mol Brain Res** **138**:273–290, 2005
137. Nakamura S, Camins MB, Hochwald GM: Pressure-absorption responses to the infusion of fluid into the spinal cord central canal of kaolin-hydrocephalic cats. **J Neurosurg** **58**:198–203, 1983
138. Nornes H, Rootwelt K, Sjaastad O: Normal pressure hydrocephalus. Long-term intracranial pressure recording. **Eur Neurol** **9**:261–274, 1973
139. Oi S, Shimoda M, Shibata M, Honda Y, Togo K, Shinoda M, et al: Pathophysiology of long-standing overt ventriculomegaly in adults. **J Neurosurg** **92**:933–940, 2000
140. Oi S, Yamada H, Sato O, Matsumoto S: Experimental models of congenital hydrocephalus and comparable clinical problems in the fetal and neonatal periods. **Childs Nerv Syst** **12**:292–302, 1996
141. Ojemann RG, Fisher CM, Adams RD, Sweet WH, New PF: Further experience with the syndrome of “normal” pressure hydrocephalus. **J Neurosurg** **31**:279–294, 1969
142. Owen-Lynch PJ, Draper CE, Mashayekhi F, Bannister CM, Miyan JA: Defective cell cycle control underlies abnormal cortical development in the hydrocephalic Texas rat. **Brain** **126**:623–631, 2003
143. Owlser BK, Pena A, Momjian S, Czosnyka Z, Czosnyka M, Harris NG, et al: Changes in cerebral blood flow during cerebrospinal fluid pressure manipulation in patients with normal pressure hydrocephalus: a methodological study. **J Cereb Blood Flow Metab** **24**:487–494, 2004
144. Patwardhan RV, Nanda A: Implanted ventricular shunts in the United States: the billion-dollar-a-year cost of hydrocephalus treatment. **Neurosurgery** **56**:139–145, 2005
145. Peña A, Bolton MD, Whitehouse H, Pickard JD: Effects of brain ventricular shape on periventricular biomechanics: a finite-element analysis. **Neurosurgery** **45**:107–118, 1999
146. Peña A, Harris NG, Bolton MD, Czosnyka M, Pickard JD: Communicating hydrocephalus: the biomechanics of progressive ventricular enlargement revisited. **Acta Neurochir Suppl** **81**:59–63, 2002
147. Pople IK, Bayston R, Hayward RD: Infection of cerebrospinal fluid shunts in infants: a study of etiological factors. **J Neurosurg** **77**:29–36, 1992
148. Portnoy HD: Electroconvulsive activity in SAH. **J Neurosurg** **53**:272–273, 1980
149. Raftopoulos C, Deleval J, Chaskis C, Leonard A, Cantraine F, Desmyttere F, et al: Cognitive recovery in idiopathic normal pressure hydrocephalus: a prospective study. **Neurosurgery** **35**:397–405, 1994
150. Ragel BT, Browd SR, Schmidt RH: Surgical shunt infection: significant reduction when using intraventricular and systemic antibiotic agents. **J Neurosurg** **105**:242–247, 2006
151. Raimondi AJ, Bailey OT, McLone DG, Lawson RF, Echeverry A: The pathophysiology and morphology of murine hydrocephalus in Hy-3 and Ch mutants. **Surg Neurol** **1**:50–55, 1973
152. Relkin N, Marmarou A, Klinge P, Bergsneider M, Black PM: Diagnosing idiopathic normal-pressure hydrocephalus. **Neurosurgery** **57** (3 Suppl):S4–S16, 2005
153. Richards HK, Bucknall RM, Jones HC, Pickard JD: Uncoupling

Priorities for hydrocephalus research

- of LCBF and LCGU in two different models of hydrocephalus: a review. **Childs Nerv Syst** 11:288–292, 1995
154. Robinson ML, Allen CE, Davy BE, Durfee WJ, Elder FF, Elliott CS, et al: Genetic mapping of an insertional hydrocephalus-inducing mutation allelic to *hy3*. **Mamm Genome** 13:625–632, 2002
155. Rolf B, Kutsche M, Bartsch U: Severe hydrocephalus in L1-deficient mice. **Brain Res** 891:247–252, 2001
156. Sato O, Takei F, Yamada S: Hydrocephalus: is impaired cerebrospinal fluid circulation only one problem involved? **Childs Nerv Syst** 10:151–155, 1994
157. Savolainen S, Hurskainen H, Paljärvi L, Alafuzoff I, Vapalahti M: Five-year outcome of normal pressure hydrocephalus with or without a shunt: predictive value of the clinical signs, neuropsychological evaluation and infusion test. **Acta Neurochir (Wien)** 144:515–523, 2002
158. Schmidt B, Czosnyka M, Schwarze JJ, Sander D, Gerstner W, Lumenta CB, et al: Evaluation of a method for noninvasive intracranial pressure assessment during infusion studies in patients with hydrocephalus. **J Neurosurg** 92:793–800, 2000
159. Schrandt-Stumpel C, Fryns JP: Congenital hydrocephalus: nomenclature and guidelines for clinical approach and genetic counselling. **Eur J Pediatr** 157:355–362, 1998
160. Schuhmann MU, Ostrowski KR, Draper EJ, Chu JW, Ham SD, Sood S, et al: The value of C-reactive protein in the management of shunt infections. **J Neurosurg** 103 (3 Suppl):223–230, 2005
161. Sciubba DM, Stuart RM, McGirt MJ, Woodworth GF, Samdani A, Carson B, et al: Effect of antibiotic-impregnated shunt catheters in decreasing the incidence of shunt infection in the treatment of hydrocephalus. **J Neurosurg** 103 (2 Suppl):131–136, 2005
162. Shiino A, Nishida Y, Yasuda H, Suzuki M, Matsuda M, Inubushi T: Magnetic resonance spectroscopic determination of a neuronal and axonal marker in white matter predicts reversibility of deficits in secondary normal pressure hydrocephalus. **J Neurol Neurosurg Psychiatry** 75:1141–1148, 2004
163. Silverberg GD, Mayo M, Saul T, Carvalho J, McGuire D: Novel ventriculo-peritoneal shunt in Alzheimer's disease cerebrospinal fluid biomarkers. **Expert Rev Neurother** 4:97–107, 2004
164. Sorteberg A, Eide PK, Fremming AD: A prospective study on the clinical effect of surgical treatment of normal pressure hydrocephalus: the value of hydrodynamic evaluation. **Br J Neurosurg** 18:149–157, 2004
165. Stephensen H, Tisell M, Wikkelsø C: There is no transmante pressure gradient in communicating or noncommunicating hydrocephalus. **Neurosurgery** 50:763–771, 2002
166. Stopa EG, Berzin TM, Kim S, Song P, Kuo-LeBlank V, Rodriguez-Wolf M, et al: Human choroid plexus growth factors: what are the implications for CSF dynamics in Alzheimer's disease? **Exp Neurol** 167:40–47, 2001
167. Sullivan PF, Keefe RS, Lange LA, Lange EM, Stroup TS, Lieberman J, et al: NCAM1 and neurocognition in schizophrenia. **Biol Psychiatry** 61:902–910, 2007
168. Tada T, Kanaji M, Kobayashi S: Induction of communicating hydrocephalus in mice by intrathecal injection of human recombinant transforming growth factor- β 1. **J Neuroimmunol** 50:153–158, 1994
169. Tada T, Zhan H, Tanaka Y, Hongo K, Matsumoto K, Nakamura T: Intraventricular administration of hepatocyte growth factor treats mouse communicating hydrocephalus induced by transforming growth factor β 1. **Neurobiol Dis** 21:576–586, 2006
170. Tedeschi E, Hasselbalch SG, Waldemar G, Juhler M, Høgh P, Holm S, et al: Heterogeneous cerebral glucose metabolism in normal pressure hydrocephalus. **J Neurol Neurosurg Psychiatry** 59:608–615, 1995
171. Tenti G, Drake JM, Sivaloganathan S: Brain biomechanics: mathematical modeling of hydrocephalus. **Neurol Res** 22:19–24, 2000
172. Thomas G, McGirt MJ, Woodworth G, Heidler J, Rigamonti D, Hillis AE, et al: Baseline neuropsychological profile and cognitive response to cerebrospinal fluid shunting for idiopathic normal pressure hydrocephalus. **Dement Geriatr Cogn Disord** 20:163–168, 2005
173. Tisell M, Edsbacke M, Stephensen H, Czosnyka M, Wikkelsø C: Elastance correlates with outcome after endoscopic third ventriculostomy in adults with hydrocephalus caused by primary aqueductal stenosis. **Neurosurgery** 50:70–77, 2002
174. Ulfing N, Bohl J, Neudörfer F, Rezaie P: Brain macrophages and microglia in human fetal hydrocephalus. **Brain Dev** 26:307–315, 2004
175. Ulfing N, Szabo A, Bohl J: Effect of fetal hydrocephalus on the distribution patterns of calcium-binding proteins in the human occipital cortex. **Pediatr Neurosurg** 34:20–32, 2001
176. Ursino M: A mathematical study of human intracranial hydrodynamics. Part 2—Simulation of clinical tests. **Ann Biomed Eng** 16:403–416, 1988
177. Van Camp G, Vits L, Coucke P, Lyonnet S, Schrandt-Stumpel C, Darby J, et al: A duplication in the L1CAM gene associated with X-linked hydrocephalus. **Nat Genet** 4:421–425, 1993
178. Wagner C, Batiz LF, Rodriguez S, Jiménez AJ, Pérez P, Tomé M, et al: Cellular mechanisms involved in the stenosis and obliteration of the cerebral aqueduct of *hyh* mutant mice developing congenital hydrocephalus. **J Neuropathol Exp Neurol** 62:1019–1040, 2003
179. Wagshul ME, Chen JJ, Egnor MR, McCormack EJ, Roche PE: Amplitude and phase of cerebrospinal fluid pulsations: experimental studies and review of the literature. **J Neurosurg** 104:810–819, 2006
180. Wagshul ME, Egnor MR, McCormack EJ, Roche PE: Stroke volume ratio as a measure of intracranial flow dysfunction. **Proc Int Soc Magn Reson Med** 1241, 2005
181. Waldemar G, Schmidt JF, Delecluse F, Andersen AR, Gjerris F, Paulson OB: High resolution SPECT with [^{99m}Tc]-d, l-HMPAO in normal pressure hydrocephalus before and after shunt operation. **J Neurol Neurosurg Psychiatry** 56:655–664, 1993
182. Whitehead WE, Kestle JR: The treatment of cerebrospinal fluid shunt infections. Results from a practice survey of the American Society of Pediatric Neurosurgeons. **Pediatr Neurosurg** 35:205–210, 2001
183. Whitelaw A, Christie S, Pople I: Transforming growth factor- β 1: a possible signal molecule for posthemorrhagic hydrocephalus? **Pediatr Res** 46:576–580, 1999
184. Williams MA, Sharkey P, Van Doren D, Thomas G, Rigamonti D: Influence of shunt surgery on health care expenditures of elderly fee-for-service Medicare beneficiaries with hydrocephalus. **J Neurosurg** 107:21–28, 2007
185. Wilson RK, Williams MA: Evidence that congenital hydrocephalus is a precursor to idiopathic normal pressure hydrocephalus in only a subset of patients. **J Neurol Neurosurg Psychiatry** 78:508–511, 2007
186. Young HF, Nulsen FE, Weiss MH, Thomas P: The relationship of intelligence and cerebral mantle in treated infantile hydrocephalus. (IQ potential in hydrocephalic children.) **Pediatrics** 52:38–44, 1973
187. Zervas NT, Cosman ER, Cosman BJ: A pressure-balanced radiotelemetry system for the measurement of intracranial pressure. A preliminary design report. **J Neurosurg** 47:899–911, 1977
188. Zhang J, Williams MA, Rigamonti D: Genetics of human hydrocephalus. **J Neurol** 253:1255–1266, 2006
189. Zonta M, Angulo MC, Gobbo S, Rosengarten B, Hossmann KA, Pozzan T, et al: Neuron-to-astrocyte signaling is central to the dynamic control of brain microcirculation. **Nat Neurosci** 6:43–50, 2003

Manuscript submitted March 29, 2007.

Accepted July 11, 2007.

Address correspondence to: James P. McAllister, Ph.D., Department of Neurosurgery, Division of Pediatric Neurosurgery, Primary Children's Medical Center, 100 North Medical Drive, Suite 1475, Salt Lake City, Utah 84113-1130. email: pat.mcallister@hsc.utah.edu.