

Guanylyl Cyclase Activating Protein

A CALCIUM-SENSITIVE REGULATOR OF PHOTOTRANSDUCTION*

(Received for publication, May 9, 1995, and in revised form, July 19, 1995)

Wojciech A. Gorczyca[‡], Arthur S. Polans[§], Irina G. Surgucheva[¶], Iswari Subbaraya[¶],
Wolfgang Baehr[¶], and Krzysztof Palczewski^{†***}

From the Departments of [†]Ophthalmology and ^{**}Pharmacology, University of Washington, Seattle, Washington 98195, the [§]R. S. Dow Neurological Sciences Institute, Good Samaritan Hospital, Portland, Oregon 97209, and the [¶]Department of Ophthalmology, Baylor College of Medicine, Houston, Texas 77030

Guanylyl cyclase activating protein (GCAP1) has been proposed to act as a calcium-dependent regulator of retinal photoreceptor guanylyl cyclase (GC) activity. Using immunocytochemical and biochemical methods, we show here that GCAP1 is present in rod and cone photoreceptor outer segments where phototransduction occurs. Recombinant and native GCAP1 activate recombinant human retGC (outer segment-specific GC) and endogenous GC(s) in rod outer segment (ROS) membranes at low calcium. In addition, we isolate and clone a retinal homolog, termed GCAP2, that shows ~50% identity with GCAP1. Like GCAP1, GCAP2 activates photoreceptor GC in a calcium-dependent manner. Both GCAP1 and GCAP2 presumably act on GCs by a similar mechanism; however, GCAP1 specifically localizes to photoreceptor outer segments, while in these experiments GCAP2 was isolated from extracts of retina but not ROS. These results demonstrate that GCAP1 is an activator of ROS GC, while the finding of a second activator, GCAP2, suggests that a similar mechanism of GC regulation may be present in outer segments, other subcellular compartments of the photoreceptor, or other cell types.

In vertebrate photoreceptor cells, the synthesis and hydrolysis of cyclic GMP (cGMP) are critical steps in phototransduction. In response to light, a cascade of reactions in the photoreceptor outer segment leads to the hydrolysis of cGMP and the closure of cGMP-gated cation channels in the outer segment plasma membrane. As a consequence, there is a reduction in the amount of calcium entering the cell. Calcium efflux owing to the Na⁺:K⁺, Ca²⁺ exchanger, however, is unaffected by light, resulting in a net decrease in the concentration of internal free calcium. This decrease in the calcium concentration leads to the activation of guanylyl cyclase (GC), which in part restores the dark conditions of the photoreceptor cell (reviewed

by Lagnado and Baylor (1992)).

Photoreceptor GC, a member of the particulate GC family (Koch, 1991; Shyjan *et al.*, 1992; Goraczniak *et al.*, 1994; Umbarger *et al.*, 1992; Dizhoor *et al.*, 1994; Liu *et al.*, 1994), responds to an activator that senses changes in the calcium concentration (Lolley and Racz, 1982; Koch and Stryer, 1988). Recently, we proposed that a calcium binding protein isolated from rod outer segments (ROS),¹ guanylyl cyclase activating protein (GCAP, termed here GCAP1), mediates this process (Gorczyca *et al.*, 1994a), and its molecular properties were described (Palczewski *et al.*, 1994). GCAP1 restores the calcium sensitivity of GC in a reconstituted system, and it decreases the sensitivity, time-to-peak, and recovery time of the light response following its introduction into intact ROS (Gorczyca *et al.*, 1994a). The molecular cloning of GCAP1 from bovine, human, mouse, frog (Palczewski *et al.*, 1994), and chicken² retina and the genomic organization of mouse and human GCAP1 (Subbaraya *et al.*, 1994) demonstrate strong sequence conservation between species, conservation of three putative calcium binding loops, and relatedness to other neuronal calcium-binding proteins of the calmodulin superfamily. Transcripts encoding GCAP1 were localized to photoreceptor cells by *in situ* hybridization (Palczewski *et al.*, 1994; Subbaraya *et al.*, 1994), but the precise localization of the protein is not known. Independently, Dizhoor *et al.* (1994) proposed that another protein, p24, was responsible for the calcium sensitivity of photoreceptor GC.

In this paper, we describe the cellular localization of bovine GCAP1 by immunocytochemical and biochemical methods and provide further evidence that GCAP1 is a key element in the activation of photoreceptor GC. In addition, we show that the retina contains a second GC activator, GCAP2, that is identical with p24 and closely related to GCAP1.

MATERIALS AND METHODS

Purification of ROS and GCAP1—Fresh bovine eyes were obtained from a local slaughterhouse, and the retinas were dissected under dim red light. ROS were prepared according to Papermaster (1982). GCAP1 was purified as described previously (Gorczyca *et al.*, 1994a).

Affinity Chromatography—A soluble extract containing GCAP1 was prepared from either bovine ROS (equivalent to 50 bovine retinas) or from 25 bovine retinas with 25 ml of water, containing 1 mM benzamide. The extract was separated from ROS or retinal particulates by centrifugation (48,000 × g for 30 min) and loaded onto an antibody-Sepharose column (mAb G-2; 6 mg of antibody per 1 ml of the CNBr-activated Sepharose; 1 × 2 cm) equilibrated with 10 mM 1,3-bis(tris(hy-

* This work was supported in part by United States Public Health Service Grants EY08061 and EY01730 (to K. P.), EY07089 (to A. S. P.), and EY08123 (to W. B.) and an award from Research to Prevent Blindness, Inc. to the Departments of Ophthalmology at the University of Washington and at Baylor College of Medicine. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.




The nucleotide sequence(s) reported in this paper has been submitted to the GenBank™/EMBL Data Bank with accession number(s) L43001.

Present address: Moran Eye Center, University of Utah Health Sciences Center, Salt Lake City, UT 84132.

‡ Recipient of a Jules and Doris Stein Research to Prevent Blindness Professorship. To whom correspondence should be addressed: Dept. of Ophthalmology, University of Washington, Box 356485, Seattle, WA 98195-6485. Tel.: 206-543-9074; Fax: 206-543-4414.

¹ The abbreviations used are: ROS, rod outer segment; BTP, 1,3-bis(tris(hydroxymethyl)methylamino)propane; HPLC, high performance liquid chromatography; PCR, polymerase chain reaction; PAGE, polyacrylamide gel electrophoresis; pAb, polyclonal antibody; mAb, monoclonal antibody; GTP_γS, guanosine 5'-O-(thiotriphosphate).

² S. Semple-Rowland and W. Baehr, unpublished results.





Downloaded from www.jbc.org at UNIV OF UTAH on August 13, 2007

Chemical Biology

1211

1212

Control of Protein-Ligand Interactions by Phosphorylation



Genomic Analysis of the *Arabidopsis thaliana* Genome


A

65

655

655

63






Fig. 1. Binding of GCAP1 to BDNF. **(A)** Purified GCAP1 (100 ng) was incubated with increasing amounts of purified BDNF. The association was detected by Western blotting, probed with IgG of the anti-GCAP1 antibody. **(B)** Purified GCAP1 (100 ng) was incubated with BDNF (100 ng) for 10 min. The association was detected by Western blotting, probed with IgG of the anti-GCAP1 antibody.

the same calcium-dependent manner as shown in Fig. 1A. The availability of immunoprecipitates of GCAP1 was not further decreasing after increasing their concentration, whereas, after immunoprecipitation, the amount of GCAP2 bound to BDNF was not detected in our Western blots. Thus, our Western blots suggest that GCAP2 in BDNF compared to other BDNF-associated proteins may be located in the plasma membrane or other parts of the retinal inner plexiform layer, but not the regulation of phototransduction at photoreceptor terminal. GCAP2 may be involved in other retinal roles and possibly, other cells and signaling systems. Alternatively, GCAP2 is different from GCAP1, it can be selectively lost during the purifying procedure of GCAP1. Therefore, we did not believe that GCAP2 is involved in the BDNF signaling pathway. Our results have not only identified the binding of GCAP1 to BDNF, but also suggest that GCAP1 is involved in BDNF signaling pathway. GCAP1 is a calcium-binding protein located in the inner plexiform layer of the retina. It is a member of the GCAP family, which is a group of calcium-binding proteins that are involved in the regulation of phototransduction. GCAP1 is a member of the GCAP family, which is a group of calcium-binding proteins that are involved in the regulation of phototransduction. GCAP1 is a member of the GCAP family, which is a group of calcium-binding proteins that are involved in the regulation of phototransduction.

- Neubert, T. A. (1994) *J. Biol. Chem.* **269**, 21067-21071
- Kawamura, S. (1993) *Nature* **362**, 855-857
- Kawamura, S., and Murakami, M. (1991) *Nature* **349**, 420-423
- Koch, K.-W. (1991) *J. Biol. Chem.* **266**, 8634-8637
- Koch, K.-W., and Stryer, L. (1988) *Nature* **334**, 64-66
- Kretsinger, R. H. (1980) *CRC Crit. Rev. Biochem.* **8**, 119-174
- Laemmli, U. K. (1970) *Nature* **227**, 680-685
- Lagnado, L., and Baylor, D. (1992) *Neuron* **8**, 995-1002
- Lambrecht, H. G., and Koch, K. W. (1991) *EMBO J.* **10**, 793-798
- Lerea, C. L., Bunt-Milam, A. H., and Hurley, J. B. (1989) *Neuron* **3**, 367-376
- Liu, X., Senoo, K., Nishizawa, Y., Hayashi, F., Yamazaki, A., Matsumoto, H., Wakabayashi, T., and Usukura, J. (1994) *Exp. Eye Res.* **59**, 761-768
- Lolley, R. N., and Racz, E. (1982) *Vision Res.* **22**, 1481-1486
- Lowe, D. G., Dizhoor, A. M., Liu, K., Gu, Q., Spencer, M., Laura, R., Lu, L., and Hurley, J. B. (1995) *Proc. Natl. Acad. Sci. U. S. A.* **92**, 5535-5539
- McGinnis, J. F., Stepanik, P. L., Baehr, W., Subbaraya, I., and Lerious, V. (1992) *FEBS Lett.* **302**, 172-176
- Murakami, A., Yajima, T., and Inana, G. (1992) *Biochem. Biophys. Res. Commun.* **187**, 234-244
- Palczewska, K., Buczylo, J., Lebiada, L., Crabb, J. W., and Polans, A. S. (1993) *J. Biol. Chem.* **268**, 6004-6013
- Palczewski, K., Subbaraya, I., Gorczyca, W. A., Helekar, B. S., Ruiz, C. C., Ohguro, H., Huang, J., Zhao, X., Crabb, J. W., Johnson, R. S., Walsh, K. A., Gray-Keller, M. P., Detwiler, P. B., and Baehr, W. (1994) *Neuron* **13**, 395-404
- Papermaster, D. S. (1982) *Methods Enzymol.* **81**, 48-52
- Pittler, S. J., and Baehr, W. (1991) *Proc. Natl. Acad. Sci. U. S. A.* **88**, 8322-8326
- Polans, A. S., and Burton, M. D. (1988) *Invest. Ophthalmol. Vis. Sci.* **29**, 1523-1532
- Polans, A. S., Buczylo, J., Crabb, J., and Palczewski, K. (1991) *J. Cell Biol.* **112**, 981-989
- Qin, N., and Baehr, W. (1993) *FEBS Lett.* **321**, 6-10
- Qin, N., and Baehr, W. (1994) *J. Biol. Chem.* **269**, 3265-3271
- Rieke, F., and Schwartz, E. A. (1994) *Neuron* **13**, 863-873
- Schoenmakers, T. J., Visser, G. J., Plik, G., and Theuvsen, A. P. (1992) *BioTechniques* **12**, 870-874
- Shyjan, A. W., de Sauvage, F. J., Gillett, N. A., Goeddel, D. V., and Lowe, D. G. (1992) *Neuron* **9**, 727-737
- Strynadka, N. C. J., and James, M. N. G. (1989) *Annu. Rev. Biochem.* **58**, 951-998
- Subbaraya, I., Ruiz, C. C., Helekar, B. S., Zhao, X., Gorczyca, W. A., Pettenati, M. J., Rao, P. N., Palczewski, K., and Baehr, W. (1994) *J. Biol. Chem.* **269**, 31080-31089
- Umbarger, K. O., Yamazaki, M., Hutson, L. D., Hayashi, F., and Yamazaki, A. (1992) *J. Biol. Chem.* **267**, 19494-19502
- Yang, R.-B., Foster, D. C., Garbers, D. L., and Fulle, H.-J. (1995) *Proc. Natl. Acad. Sci. U. S. A.* **92**, 602-606